

RESEARCH ARTICLE

Resource Seasonality and Reproduction Predict Fission–Fusion Dynamics in Black-and-White Ruffed Lemurs (*Varecia variegata*)ANDREA L. BADEN^{1,2,3*}, TIMOTHY H. WEBSTER⁴, AND JASON M. KAMILAR^{5,6,7}¹Department of Anthropology, Hunter College, New York, New York²Graduate Center, City University of New York, New York, New York³New York Consortium in Evolutionary Primatology (NYCEP), New York, New York⁴Department of Anthropology, Yale University, New Haven, Connecticut⁵Department of Anthropology, University of Massachusetts, Amherst, Massachusetts⁶Graduate Program in Organismic and Evolutionary Biology, University of Massachusetts, Amherst, Massachusetts⁷School of Human Evolution and Social Change, Arizona State University, Tempe, Arizona

Ruffed lemurs (genus *Varecia*) are often described as having a flexible social organization, such that both cohesive (low fission–fusion dynamics) and fluid (high fission–fusion dynamics) grouping patterns have been observed. In ruffed lemur communities with high fission–fusion dynamics, group members vary in their temporal and spatial dispersion throughout a communally defended territory. These patterns have been likened to those observed in several haplorrhine species that exhibit the most fluid types of fission–fusion social organization (e.g., *Pan* and *Ateles*). To substantiate and further refine these claims, we describe the fission–fusion dynamics of a black-and-white ruffed lemur (*Varecia variegata*) community at Mangevo, an undisturbed primary rainforest site in Ranomafana National Park, Madagascar. We collected instantaneous group scan samples from August 2007–December 2008 (4,044 observation hours) to study and characterize patterns of subgroup size, composition, cohesion, and social association. In 16 consecutive months, we never found all members of the community together. In fact, individuals spent nearly half of their time alone. Subgroups were small, cohesive, and typically of mixed-sex composition. Mixed-sex subgroups were significantly larger, less cohesive, and more common than either male-only or female-only subgroups. Subgroup dynamics were related to shifts in climate, phenology of preferred fruit species, and female reproductive state. On average, association indices were low. Males and females were equally gregarious; however, adult male–male associations were significantly weaker than any other association type. Results presented herein document striking differences in fission–fusion dynamics between black-and-white ruffed lemurs and haplorrhines, while also demonstrating many broad-scale similarities to haplorrhine taxa that possess the most fluid fission–fusion societies. *Am. J. Primatol.* 78:256–279, 2016. © 2015 Wiley Periodicals, Inc.

Key words: association; subgroup size; party type; cohesion; seasonality; Madagascar

INTRODUCTION

Gregariousness affords both costs and benefits to group-living primates [Alexander, 1974; Majolo et al., 2008; van Schaik, 1989]. Group size, in particular, is considered among the most important factors modulating behavior and individual fitness [Janson & Goldsmith, 1995; Koenig, 2002; Sterck et al., 1997; van Schaik, 1983; Wrangham, 1980]. As group size increases, individuals may benefit from improved success in intergroup contests over access to food resources [Janson & van Schaik, 1988; van Schaik, 1989; Wrangham, 1980], and reduced predation pressure [Hill & Lee, 1998; Sterck et al., 1997; van Schaik, 1983]; however, they may simultaneously face greater intragroup feeding competition [Alexander, 1974; Clutton-Brock & Harvey, 1977; Isbell, 1991; Janson & van Schaik, 1988; Terborgh &

Contract grant sponsor: National Science Foundation; contract grant number: DDIG BSC-0725975; contract grant sponsor: US Department of State Fulbright Fellowship; contract grant sponsor: The Leakey Foundation; contract grant sponsor: Primate Conservation, Inc.; contract grant sponsor: Conservation International's Primate Action Fund

*Correspondence to: Andrea L. Baden, Department of Anthropology, Hunter College 695 Park Avenue, New York, NY 10065. E-mail: andrea.baden@hunter.cuny.edu

Received 25 November 2014; revised 4 October 2015; revision accepted 30 October 2015

DOI: 10.1002/ajp.22507

Published online 25 November 2015 in Wiley Online Library (wileyonlinelibrary.com).

Janson, 1986; van Schaik, 1989], increased conspicuousness to predators [e.g., Stanford, 1998] or to infanticidal males [e.g., Crockett and Janson, 2000], higher physiological stress [Pride, 2005; Snaith et al., 2008], and greater susceptibility to disease [Caillaud et al., 2013; Nunn & Heymann, 2005; Rifkin et al., 2012]. Group size should thus reflect an optimization of the benefits and costs of group living, primarily in relation to two key factors: resource competition (over food or mates) and predator avoidance [Terborgh & Janson, 1986]. Local selective pressures can change the benefit-to-cost ratio of gregariousness, and can influence the size and composition of social groups; in fact, optimal group size may change with time, even within the same population [van Schaik & van Hooff, 1983].

In some species, group members are able to individually optimize the costs and benefits of group living via a strategy known as “fission–fusion” [*sensu* Kummer, 1971]. In these taxa, individuals form socially and geographically circumscribed unit groups (i.e., communities) within which members associate in temporary, flexible subunits (i.e., parties or subgroups) that vary in size, cohesion, membership, and duration [e.g., Mitani et al., 2002b]. Although the precise social and ecological factors that promote fission–fusion social dynamics are not fully understood, this behavior is thought to function primarily to minimize within-group competition over access to food and mates [Chapman et al., 1995; Mitani et al., 2002a and references therein]. Specifically, subgroup size is often closely tied to the availability of high-quality food resources, particularly ripe fruit [Asensio et al., 2009; Chapman et al., 1995; Symington, 1987, 1988b and references therein]. For example, groups fission and subgroups tend to be smaller and/or less cohesive when resources are scarce [e.g., Aureli & Schaffner, 2008; Anderson et al., 2002; Boesch, 1996; Chapman et al., 1995; Doran, 1997; Ghiglieri, 1984; Klein & Klein, 1977; Matsumoto-Oda et al., 1998; Mitani et al., 2002a; Morland, 1991a; Newton-Fisher et al., 2000; Shimooka, 2003; Symington, 1988a, 1990; Vasey, 2006; Wrangham, 1977, 1980; Wrangham et al., 1992, 1996; but see Stanford et al., 1994; Boesch, 1996; Hashimoto et al., 2001, 2003], when foods are patchily distributed (i.e., non-defensible), or when patches are small [Chapman, 1990a,b; Chapman et al., 1995; de Moraes et al., 1998; Isabirye-Basuta, 1988; Klein & Klein, 1977; Newton-Fisher et al., 2000; Potts et al., 2011; Shimooka, 2003; Stevenson et al., 1998; Symington, 1988b, 1990; Vasey, 2000; White & Wrangham, 1988; Wittiger & Boesch, 2013; Wrangham, 1977, 1980, 1986].

In addition, variation in subgroup size has been attributed to demographic factors (e.g., community size, sex ratio [Boesch, 1996; Goodall, 1986; Newton-Fisher, 1999; Newton-Fisher et al., 2000]) and behavioral context (e.g., cooperative hunting or

territorial patrols [Boesch & Boesch, 1989; Wallace, 2008]), as well as infant socialization [Chapman et al., 1994; Newton-Fisher, 1999; Williams et al., 2002] and female reproductive state [Anderson et al., 2002; Chapman, 1990a; Boesch & Boesch-Acher-mann, 2000; Goodall, 1986; Hashimoto et al., 2001; Matsumoto-Oda, 1999; Matsumoto-Oda et al., 1998; Mitani et al., 2002a; Sakura, 1994; Shimooka, 2003; Stanford et al., 1994; Symington, 1987; Wrangham, 2000]. Mixed-sex subgroups are typically larger when estrus/cycling females are present. Non-cycling adult females (i.e., those that are pregnant, lactating, or post-reproductive) are less commonly observed in large bisexual parties [Matsumoto-Oda, 1999], and instead, pregnant females tend to be less social [Goodall, 1986; Nishida, 1990], especially just before parturition [Pusey et al., 2008]. Likewise, lactating females generally spend more time alone [Goodall, 1986; Hasegawa, 1990; Kawanaka, 1984; Sakura, 1994; Uehara, 1981; Wrangham & Smuts, 1980; but see Shimooka, 2003; Vasey, 2007] or in small subgroups with other lactating females [Sakura, 1994; Slater et al., 2009].

Because primate social organization has historically been classified into a handful of modal categories, only primates exhibiting the most extreme forms of fission–fusion have typically been classified as having a “fission–fusion social organization” [Aureli et al., 2008; Clutton-Brock & Harvey, 1977; Di Fiore & Rendall, 1994; Kappeler, 1997; Strier, 1994]. For this reason, our knowledge of primate fission–fusion stems primarily from studies of chimpanzees (*Pan troglodytes*) [Boesch & Boesch-Acher-mann, 2000; Mitani et al., 2002a,b; Nishida & Hiraiwa-Hasegawa, 1987], bonobos (*P. paniscus*) [Furuichi, 2009; Hohmann & Fruth, 2002; Stumpf, 2011], and spider monkeys [*Ateles* spp.: Chapman et al., 1995; Di Fiore et al., 2011; Symington, 1990]. Convergent patterns of fission–fusion among these taxa suggest that similar social systems have evolved in response to similar selection pressures in distantly related taxa. However, with increased knowledge of individual ranging patterns, it is becoming clear that flexible spatiotemporal grouping patterns in primates and other vertebrates are more common—and more complex—than generally recognized [e.g., Aureli et al., 2008; Kinzey & Cunningham, 1994; Struhsaker & Leland, 1979; Sussman & Garber, 2011]. In fact, fission–fusion dynamics [*sensu* Aureli et al., 2008] can be quantified in any primate group that varies, even minimally, in terms of its size, membership, and spatial cohesion through time (e.g., brown lemurs [Overdorff et al., 2003; Toborowsky, 2008], capuchins [Lynch-Alfaro, 2007], geladas [Snyder-Mackler et al., 2012], howler monkeys [Bezanson et al., 2008], humans [Marlowe, 2005], macaques [Ménard, 2002], murrelets [Coles et al., 2012], ring-tailed lemurs [L. Kelley, pers. comm.], saki monkeys [Veiga et al., 2006]; snub-nosed

monkeys [Ren et al., 2012], uakaris [Bowler & Bodmer, 2009; Bowler et al., 2012]).

In this study, we characterize the nature of fission–fusion dynamics in a population of black-and-white ruffed lemurs (*Varecia variegata*). Ruffed lemurs are relatively large-bodied [2.5–4.8 kg; Baden et al., 2008], arboreal frugivores endemic to the eastern rainforests of Madagascar [Baden, 2011; Balko, 1998; Morland, 1991a; Ratsimbazafy, 2002; Rigamonti, 1993; Vasey, 2000]. While initially described as living in cohesive, pair-bonded groups [Petter et al., 1977; Pollock, 1979; White, 1991], nearly all subsequent long-term studies describe ruffed lemurs as living in groups characterized by high fission–fusion social dynamics [Baden, 2011; Morland, 1991a,b; Rigamonti, 1993; Vasey, 2006; but see Balko, 1998; Britt, 1997; Ratsimbazafy, 2002]. Groups or “communities” typically range in size between 11 and 31 individuals [Baden, 2011; Morland, 1991a,b; Vasey, 2006], though members are rarely if ever all seen together in the same place at the same time. Aspects of their fission–fusion dynamics—including range use, spatial associations, and affiliative interactions—vary according to season, resource availability, and female reproductive state [Morland, 1991a,b; Rigamonti, 1993; Vasey, 2006], and their associations have been described as “female-bonded” [Morland, 1991a]. These foundational studies of fission–fusion sociality in the strepsirrhine genus *Varecia* have demonstrated some remarkable similarities with haplorrhine taxa that exhibit the most extreme forms of fission–fusion sociality (e.g., chimpanzees, bonobos, and spider monkeys). To substantiate and further refine these claims, we employ multivariate modeling to gauge the relative influence of various social and ecological factors thought to impact subgroup size and cohesion. We report on subgroup size, type, cohesion, and duration, and describe patterns of association within one communal *V. variegata* territory. Further, we evaluate whether and how ruffed lemur fission–fusion dynamics vary with seasonal shifts in climate, fruit availability, and reproductive state. Finally, to assess species and population parallels and differences, we couch our results in a larger comparative context of primate fission–fusion dynamics.

Study Site and Subjects

Data were collected from one *V. variegata* community at Mangevo bushcamp in Ranomafana National Park, Madagascar (RNP) for 16 months (August–December 2007; February–December 2008). Mangevo [21°22′60″S, 47°28′0″E] is a mid elevation (660–1,200 m) primary rainforest site within the southeastern parcel of RNP, containing 435 km² of continuous montane rainforest located in the

southeastern escarpment of Madagascar’s central high plateau [21°02′–21°25′S and 47°18′–47° 37′E; Wright et al., 2012].

All members of the Mangevo ruffed lemur community were habituated to human observers and were individually identified via radio-collars and/or unique collar-tag combinations prior to the study. Animals were collared under veterinary supervision following a strict protocol previously outlined [Glander, 1993]. At the time of the study, the community included 24 adults and subadults (8 adult females, 11 adult males, 5 subadult males). Nineteen infants were born in the 2008 birth season and were present from October to December 2008, when the study ended. Of the study subjects, 5 females and 3 males were radio-collared and targeted for regular follows. Individuals with collar-tags (but no radio-collars, $n = 16$) were opportunistically targeted for focal follows. Sampling efforts resulted in a total of 4,044 focal observation hours.

Research protocols were in compliance with and permission was granted by Stony Brook University IACUC #2005-20081449 and Madagascar’s National Parks (ANGAP/MNP), and adhered to the American Society of Primatologists (ASP) Principles for the Ethical Treatment of Non Human Primates.

Data Collection

Two teams of four observers each conducted dawn-to-dusk follows on focal individuals, (i.e., two animals were followed daily). A focal animal was located at the beginning of each observation period via radio-telemetry. Only independent individuals (adults and subadults) were targeted for follows. New focals were selected daily. Focals were never sampled on consecutive days and every effort was made to follow all focals at least once per month. If an individual with a collar-tag was located in association with a radio-collared focal individual prior to 10:00 h, this individual became the new focal for that observation period. Observation periods ranged in duration between 8 and 11 hr depending on seasonal differences in day length and time needed to locate animals at dawn.

Upon initial contact with the focal individual, we recorded the number and identities of all other individuals present within the subgroup. To do so, one observer remained with the focal individual while the remaining three team members spread out to locate and identify all other members of the subgroup. Subgroups were defined as all independent individuals (i.e., adults and subadults) within a 50 m radius of the approximate subgroup center that exhibited coordinated behavior and travel (i.e., traveling in the same direction). This threshold was chosen because: (i) four observers could reliably detect and differentiate animals within 50 m and (ii) subgroup members within this radius were assumed

to be in visual contact with each other as individuals visible to the observer were presumably also visible to each other [Lehmann & Boesch, 2004]. Our subgroup definition is similar to that described by Coles et al. [2012], in that it combined both a *general* spatial criterion [i.e., all individuals in sight: Boesch, 1996; Chapman, 1990a; Itoh & Nishida, 2007] and a *specific* spatial criterion [i.e., all individuals within a defined radius: Morgan, 2007; Newton-Fisher et al., 2000]. We also noted, *ad libitum*, the number, identity, and distance of individuals that were observed outside of the 50 m radius and monitored their individual behavior (e.g., whether the animal approached, retreated from, or exhibited coordinated travel with the subgroup being studied) while in sight. These instances were rare, and generally comprised a solitary individual traveling past the subgroup at distances of at least 100 m. However, in cases where animals spent considerable time (i.e., ≥ 30 min) traveling in the same direction as and/or coordinating activity with the subgroup being followed, these animals were, *post hoc*, considered part of the subgroup, and were included in estimates of subgroup size, composition, and cohesion.

After initial contact, we monitored subsequent changes in subgroup size, composition (age/sex class, individual identity), and cohesion (i.e., the greatest distance between any two subgroup members), as well as activity state of the focal subject using instantaneous scan sampling techniques collected at 5 min intervals [Altmann, 1974]. We recorded five activity states: (1) feed/forage (to search for, procure, ingest, or chew any food item); (2) travel (movement from one location to another, excluding changing position within a patch during feeding/foraging); (3) rest (stationary with no apparent activity); (4) social (active interaction between ≥ 2 individuals); and (5) other (e.g., self-groom). If a social interaction occurred during any of the other four activity categories (e.g., an agonistic interaction during a feeding bout), “social” was recorded. Subgroup type (i.e., the sex composition of subgroups: female only, male only, mixed sex) was assigned *post hoc* according to the composition of subgroups at each 5 min scan. Subgroup fission events were recorded when ≥ 1 individual(s) were absent from more than two consecutive group scans; fusion events were recorded when ≥ 1 individual(s) not previously belonging to the subgroup were found in association with another subgroup member during a group scan. With these methods, we recorded a total of 40,840 group scans and 2,514 changes in subgroup size and/or composition (fission and/or fusion events).

Because phenological patterns of fruit abundance and scarcity in Madagascar often closely correlate with climatic changes in temperature and rainfall [Dewar & Richard, 2007; Hemingway, 1996, 1998; Meyers & Wright, 1993; Overdorff, 1993], we collected daily measurements of total rainfall (mm),

and mean, minimum, and maximum temperatures (C°). Throughout the study, we also monitored fruiting productivity and female reproductive state as these have previously been related to patterns of fission–fusion dynamics. Figure 1 illustrates the correspondence between climate, fruit availability, and female reproductive state during the study period. Because females bore young during the birth season in late 2008, but not in 2007, data were collected during early lactation (October–December 2008), but not during mid to late lactation (January–February/March 2008).

We estimated fruit availability from 637 trees and lianas found within 12 botanical plots ($10 \times 50 \text{ m}^2$ or 500 m^2) established evenly throughout the communal home range (87.8 ha; [Baden, 2011; Baden & Gerber, in prep]). We monitored only those species known to comprise the top 25 preferred *Varecia* food resources [as per Baden & Gerber, in prep; Balko, 1998]. Within each plot, we identified the following information for each stem ≥ 5 cm diameter at breast height (DBH): species identity, position (x, y) relative to plot origin, DBH, height, crown height, and crown diameter. Plants were monitored during one week per month. Each tree was assigned a percentage score for each of four phenophases (flower buds, flowers, unripe fruit, ripe fruit), representing the estimated crown coverage. Because all four phenophases occupy the same physical space within the crown, the sum of the four scores did not exceed 100%. From these records, we calculated an estimate of monthly fruit availability (%) by summing the percent ripe and unripe fruits for each tree and then averaging across species.

Female reproductive state was noted at the beginning of each observation period. Ruffed lemurs are strict seasonal breeders and exhibit clear signs of behavioral and vaginal estrus [Brockman et al., 1987; Foerg, 1982; Morland, 1993; Pereira et al., 1988]. Thus, we were able to easily assess female reproductive state as one of three broad reproductive categories during our study: anestrus, gestation, and lactation.

Data Analyses

Characterizing ruffed lemur fission–fusion dynamics

From the full dataset (40,840 scans), we calculated descriptive statistics related to fission–fusion social dynamics within the community, including the rate of subgroup fissions and fusions, subgroup size, and subgroup cohesion. To make our data comparable to those from other studies and taxa, we present subgroup size as both absolute and relative measures (i.e., subgroup size expressed as a percentage of total group size *sensu* Boesch [1996]). We excluded the first scan in each observation period when calculating rates of fissions and fusions, as




Fig. 1. Correspondence of climate, fruit availability, and reproductive state during the study period. Dashed line represents rainfall; solid line represents temperature. Note: no infants were born in 2007, thus January–June 2008 were considered non-reproductive months. In other years, some of these months would be a continuation of lactation.

subgroup size and composition prior to the first scan was unknown.

Sampling was biased toward radio-collared females. We, therefore, analyzed only a subset of our data to ascertain subgroup type and association patterns. Scans containing unknown individuals (i.e., animals with neither radio-collars nor collar-tags) were removed and data were then divided by month according to whether the focal individual was male or female. We then randomized the point scans and selected X scans to include in the dataset for each sex, where X is equal to 90% of the point scans for the sex with the fewest scans in a given month. For example, in April 2008, there were 2,098 female focal scans and 789 male focal scans. We, therefore, randomly selected 1,420 point scans from April, such that we included 710 focal male scans (90% of 789, the smaller number) and 710 focal female scans in the subsampled dataset. This procedure resulted in a total of 11,784 point scans for inclusion in our analysis targeting equal numbers of males and females in each month. From this subsampled dataset, we calculated descriptive statistics for subgroup size, type, and sex ratio. We report means \pm standard deviations.

Characterizing patterns of association

We used Socprog 2.5 [Whitehead, 2009] to calculate association indices between pairs of individuals using the “simple ratio” index, which is appropriate if individuals are equally likely to be

correctly identified [Cairns & Schwager, 1987; Ginsberg & Young, 1992; Whitehead, 2008]. The simple ratio index is defined as

$$\frac{x}{x + y_{AB} + y_A + y_B}, \quad (1)$$

where x = the number of sampling periods in which individuals A and B were observed/associated within a given subgroup, y_{AB} = the number of sampling periods in which A and B were identified but not associated, y_A = the number of sampling periods in which only A was identified, and y_B = the number of sampling periods in which only B was identified. Because subgroup composition changed, on average, every 90 min, we set sampling periods to 6 hr to reduce autocorrelation among periods. We removed individuals observed in fewer than 10 sampling periods.

To test for sex differences in association, we performed a Mantel test implemented in Socprog 2.5 [Whitehead, 2009]. The Mantel test correlates the matrix of association indices and a 1:0 matrix indicating whether or not members of a pair are the same sex, and compares that value to a distribution of values calculated by randomly permuting the sexes of individuals, allowing the calculation of statistical significance [Mantel, 1967; Schnell et al. 1985; Whitehead, 2008]. We increased the number of permutations until P values stabilized ($n = 10,000$). Tests were two-tailed and alpha was set at 0.05.

Factors predicting subgroup size and cohesion

We constructed linear mixed models (LMM) and generalized linear mixed models (GLMM) to assess factors influencing subgroup cohesion and subgroup size, respectively. We performed two analyses for each of these dependent variables. One model contained two climate variables as predictors, rainfall and temperature (as a proxy for fruit availability). These climate variables were the first two components generated from a principal components analysis (PCA). The PCA was based on a covariation matrix and was performed using the `princomp` function in R [R Development Core Team, 2014]. The second model contained fruit availability as a predictor. In addition, all models included the following fixed effects as predictor variables: female reproductive state, activity, and subgroup type. Models assessing subgroup cohesion also included subgroup size as a predictor as we were interested in subgroup cohesion patterns independent of subgroup size. Each model designated the focal individual and data collection day as random effects. We chose these two random effects because we used our point scans as samples in the analyses. Therefore, including these variables as random effects in the models allowed us to account for repeatedly measuring the same individual and the same day. This controlled for individual level variation, as well as uneven sampling among individuals. Similarly, including data collection day as a random effect accounted for multiple scans on the same day, many of which were associated with the same values for some variables (e.g., fruit availability, daily climate). We used the `lmer` and `glmer` functions in the R package `lme4` [Bates et al., 2014] to conduct the LMM and GLMM models, respectively. For the GLMM models, “family” was set to `poisson`. The `lmerTest` package

[Kuznetsova et al., 2014] was used to produce *P* values.

For each of the four full models, we used Akaike’s Information Criterion corrected for small sample size (AICc) [Burnham & Anderson, 2002] to judge the combination of predictor variables that best fit the data. The model associated with the lowest AICc score was considered the best and other models within 2 AICc values were considered equally good [Burnham & Anderson, 2002]. This approach has several advantages over stepwise procedures [see Garamszegi, 2011 for a recent review]. Finally, we calculated the sum of AICc weights for each fixed effect to assess their relative importance. This metric varies from zero to one and fixed effects with higher values are more frequently present in the best models. Therefore, these predictors better explain the dependent variable. All AICc related metrics were calculated with the `dredge` function in the `MuMin` package for R [Bartoń, 2014].

RESULTS**Ruffed Lemur Fission–Fusion Dynamics**

We never found all community members together in the same place at the same time; indeed, we never observed more than one-third of community members together in a subgroup. Focal individuals were solitary for nearly one-half (48%) of observations. During scans where at least two individuals were present, subgroups primarily comprised two or three individuals (mean = 2.54 ± 0.85 s.d.; range = 2–7; $n = 21,219$; Fig. 2). Relative subgroup size within the community was equally low (mean = 10.59%; range = 8.33–29.17%).

Subgroups were typically cohesive, with an average group spread of $8.02 \text{ m} \pm 7.21$ s.d.; however,




Fig. 2. Frequency distribution of ruffed lemur subgroup sizes from 5 min time point scans ($n = 40,840$).




Fig. 3. Subgroup cohesion by activity type, as calculated from 5 min time point scans ($n = 40,840$).

group cohesion varied by activity type (range = 0–60 m; Fig. 3). Subgroups were most cohesive when members were engaged in social behaviors ($\text{mean}_{\text{SOC}} = 6.22 \text{ m} \pm 9.19 \text{ s.d.}$), followed by resting ($\text{mean}_{\text{RST}} = 7.70 \text{ m} \pm 7.17 \text{ s.d.}$), and feeding ($\text{mean}_{\text{FD}} = 8.24 \text{ m} \pm 6.96 \text{ s.d.}$). Subgroups were least cohesive during travel ($\text{mean}_{\text{TRV}} = 9.78 \text{ m} \pm 7.4 \text{ s.d.}$).

Subgroup size varied by subgroup type (Table I). When at least two individuals were present, mixed-sex subgroups were predominant (79.5%), followed by male-only (12.1%) and female-only (8.4%) subgroups. Mixed-sex subgroups were larger ($2.57 \text{ ind.} \pm 0.82$) than male-only ($2.25 \text{ ind.} \pm 0.43$) and female-only ($2.08 \text{ ind.} \pm 0.28$) subgroups (Table I), and sex ratios within mixed-sex subgroups were nearly equal (0.49 ± 0.38). Mixed-sex subgroups also reached larger maximum sizes than same-sex subgroups (Table I).

Males were only slightly more likely than females to be found in subgroups with at least one other individual (71% vs. 68%, respectively), and males and females were equally likely to be found in mixed-sex subgroups (62% vs. 61%, respectively).

On average, subgroup size and composition changed every 90 min. Fissions occurred every 173 min and fusions occurred every 174 min.

Social Association Patterns

Dyadic associations within the Mangevo community were generally weak, with values ranging between 0 (never associated) and 0.80 (11,765 scans; mean association index = 0.05 ± 0.03). When considering dyadic relationships among all independent community members (i.e., among adults and subadults), association indices did not differ significantly for same sex or mixed-sex dyads (Mantel test: $r = -0.08$, $P = 0.09$). However, subadult males exhibited unusually high association indices (mean = 0.09) and we noted that the association between two subadult males in particular—the highest observed association index, at 0.80—was likely inflating male association indices. We, therefore, reran analyses using adults only (11,171 records) and found adult male–male dyads exhibited significantly weaker association indices than adult female–female or mixed-sex dyads (Mantel test, $r = -0.15$, $P = 0.016$; mean adult male–male AI = 0.01 ± 0.01 ; mean adult male–female AI = 0.06 ± 0.04 ; mean adult female–male AI = 0.06 ± 0.03 ; mean adult female–female AI = 0.05 ± 0.02) (Fig. 4).

TABLE I. Variation in *Varecia variegata* Subgroup Size by Subgroup Type (Subsampled Dataset)

Sex composition	Scans (n)	% observations	Mean (\pm SD)	Min.	Max.
Solitary male	2,440	20.7	–	1	1
Solitary female	2,768	23.5	–	1	1
Males	796	6.8	1.31 ± 0.58	2	3
Females	553	4.7	1.18 ± 0.42	2	4
Mixed-sex	5,227	44.4	2.57 ± 0.82	2	6
Total	11,784				




Fig. 4. Association index values (means \pm s.d.) according to subgroup type. Dark bars represent values for adults and subadults combined, while gray bars represent values for adults only. *** $P < 0.001$

Factors Predicting Subgroup Size and Cohesion

For the GLMM predicting subgroup size, which used climate variables as a proxy for fruit availability, a single model had the lowest AICc value. No other model was within 2 AIC values of this best model. This model contained all the predictors except for activity (Table II). Based on the sum of AICc weights, subgroup type and reproductive state were the most important predictors (values of 1.00

and 0.94, respectively) (Table III). The two climate variables exhibited moderately high values (0.83 and 0.75, respectively), and activity displayed a low value (0.16). This latter result suggests that subgroup size was only weakly related to activity. Smaller subgroups (solitary individuals) were associated with the early lactation period, as well as lower temperature and rainfall values (as loaded on the climate PCA axes). Larger subgroups were associated with mixed sex subgroups (Table I).

TABLE II. Results of Generalized Linear Mixed Model Predicting Subgroup Size Using Climate Variables as a Proxy for Fruit Availability. Full Model Results Are Presented Here

Fixed effect	Estimate	Std. error	z value	P
(Intercept)	0.756	0.041	18.66	<0.001
Climate_PCA1	-0.020	0.008	-2.41	0.016
Climate_PCA2	-0.022	0.010	-2.10	0.035
ReprodState_Lactation	-0.129	0.039	-3.34	0.001
ReprodState_Nonreproductive	-0.043	0.032	-1.36	0.174
Activity_Other	0.007	0.020	0.36	0.715
Activity_Rest	-0.001	0.010	-0.09	0.931
Activity_Social	0.106	0.050	2.11	0.035
Activity_Travel	-0.006	0.017	-0.34	0.732
SubGpType_MaleOnly	-0.087	0.044	-1.98	0.048
SubGpType_MixedSex	0.222	0.017	12.75	<0.001
SubGpType_SolitaryFemale	-0.684	0.019	-36.32	<0.001
SubGpType_SolitaryMale	-0.667	0.032	-20.56	<0.001

Model AICc = 83694.7; logLikelihood = -41832.34.

The model with the best AICc value (83691.5) did not contain the activity variable.

The focal individual and data collection day were used as random effects.

Gestation is the reference category for the Reproductive State variable. Feed/forage is the reference category for the activity variable.

Female only is the reference category for the Subgroup Type variable.

The P -value associated with each listed category indicates whether it is significantly different from the reference category noted in the footer, e.g., social is associated with lower group spread than feed/forage.

TABLE III. Relative Importance of Predictor Variables for Explaining Variation in Subgroup Size and Cohesion Based on the Sum of AICc Weights

Model	Climate PCA1	Climate PCA2	Fruit availability	Reproductive state	Sex composition	Activity state	Subgroup size
Predicting subgroup size ^a	0.83	0.75	–	0.94	1	0.16	–
Predicting subgroup size ^b	–	–	1	1	1	0.06	–
Predicting subgroup cohesion ^a	0.14	0.57	–	1	1	1	1
Predicting subgroup cohesion ^b	–	–	0.98	1	1	1	1

Note: The sum of AICc weights vary from zero to one.

^aModel using climate variables as proxy for fruit availability.

^bModel using phenological plots as proxy for fruit availability.

The GLMM predicting subgroup size that included fruit availability produced similar results. A single model had the lowest AICc value and contained all the predictors except for activity (no other model was within 2 values of this best model) (Table IV). Based on the sum of AICc weights, subgroup type, reproductive state, and fruit availability were the most important predictors (all exhibited a value of 1.00) (Table III). In contrast, activity displayed a value of 0.06. Again, smaller subgroup sizes were associated with the early lactation period, whereas larger subgroup sizes were associated with mixed-sex subgroups and higher fruit availability.

Our LMM predicting subgroup cohesion and including climate variables as a proxy for fruit availability generated two best models based on AICc. The model with the lowest AICc value (105316.4) contained all predictor variables except for climate PCA 1 (Table V). An equivalent model (AICc = 105316.9) contained all predictors except for either of the climate variables. Based on the sum of

AICc weights, subgroup type, subgroup size, reproductive state, and activity were the most important predictors (all exhibited values of 1.00) (Table III). Climate PCA 2 exhibited a moderate score, 0.57, and climate PCA 1 exhibited a low score 0.14. Greater subgroup cohesion was associated with the lactation period, as well as time spent resting and social. Reduced subgroup cohesion was associated with mixed sex. Finally, subgroup size was negatively related to subgroup cohesion.

Finally, our LMM predicting subgroup cohesion and using fruit availability produced a single best model based on AICc (no other model was within 2 values of this best model). This model contained all predictor variables (Table VI). Based on the sum of AICc weights, all the predictors were nearly equally important (Table III). All predictors exhibited a sum of AICc weight value of 1.00, except for fruit availability, which was 0.98. Greater subgroup cohesion was associated with the lactation period, as well as time spent resting and social. Higher fruit availability was also associated with greater

TABLE IV. Results of Generalized Linear Mixed Model Predicting Subgroup Size Using Fruit Availability From Botanical Plots. Full Model Results Are Presented Here. The P Value Associated with Each Listed Category Indicates Whether It is Significantly Different From the Reference Category Noted in the Footer, e.g., Social is Associated With Lower Group Spread Than Feed/Forage

Fixed effect	Estimate	Std. error	z value	P
(Intercept)	0.524	0.053	9.90	<0.001
TotalFruitAvailable	2.713	0.458	5.92	<0.001
ReprodState_Lactation	–0.125	0.032	–3.96	<0.001
ReprodState_Nonreproductive	0.016	0.028	0.57	0.572
Activity_Other	0.004	0.021	0.19	0.848
Activity_Rest	–0.002	0.011	–0.14	0.888
Activity_Social	0.089	0.060	1.48	0.139
Activity_Travel	–0.007	0.018	–0.40	0.689
SubGpType_MaleOnly	–0.112	0.046	–2.45	0.014
SubGpType_MixedSex	0.217	0.018	12.09	<0.001
SubGpType_SolitaryFemale	–0.681	0.020	–34.76	<0.001
SubGpType_SolitaryMale	–0.685	0.034	–19.87	<0.001

Model AICc = 76652.4; logLikelihood = –38312.22.

The model with the best AICc value (76646.9) did not contain the activity variable.

The focal individual and data collection day were used as random effects.

Gestation is the reference category for the Reproductive State variable.

Feed/forage is the reference category for the Activity variable. Female only is the reference category for the Subgroup Type variable.

TABLE V. Results of Linear Mixed Model Predicting Subgroup Cohesion Using Climate Variables as a Proxy for Fruit Availability. Full Model Results Are Presented Here. The *P* Value Associated with Each Listed Category Indicates Whether It is Significantly Different From the Reference Category Noted in the Footer, e.g., Social is Associated With Lower Group Spread than Feed/Forage

Fixed effect	Estimate	Std. error	<i>t</i> value	<i>P</i>
(Intercept)	0.081	0.754	0.11	0.914
Climate_PCA1	0.038	0.179	0.21	0.831
Climate_PCA2	0.435	0.224	1.94	0.054
ReprodState_Lactation	−1.943	0.809	−2.40	0.017
ReprodState_Nonreproductive	1.754	0.665	2.64	0.009
Activity_Other	−0.050	0.237	−0.21	0.832
Activity_Rest	−0.699	0.115	−6.08	<0.001
Activity_Social	−3.052	0.520	−5.87	<0.001
Activity_Travel	2.508	0.201	12.50	<0.001
SubGpType_MaleOnly	2.242	0.430	5.21	<0.001
SubGpType_MixedSex	1.361	0.171	7.97	<0.001
SubGpType_SolitaryFemale	4.970	1.144	4.34	<0.001
SubGpType_SolitaryMale	5.737	2.333	2.46	0.014
Subgroup size	2.637	0.081	32.51	<0.001

Model AICc = 105319.9; logLikelihood = −52642.94.

The model with the best AICc value (105316.4) did not contain Climate_PCA1.

The focal individual and data collection day were used as random effects.

Gestation is the reference category for the Reproductive State variable.

Feed/forage is the reference category for the Activity variable.

Female only is the reference category for the Subgroup Type variable.

subgroup cohesion values, though the *P* value was 0.08 in the full model. Finally, reduced subgroup size was associated with mixed sex subgroup types. Again, subgroup size was negatively related to subgroup cohesion.

DISCUSSION

Ruffed Lemur Fission Fusion Dynamics

Ruffed lemurs are often described as having a flexible social organization, such that both cohesive

(low fission–fusion dynamics) and fluid grouping patterns (high fission–fusion dynamics) have been observed [reviewed in Vasey, 2003, 2006]. Results from this study demonstrate that, much like patterns characteristic of northern ruffed lemur populations [Morland, 1991a,b; Rigamonti, 1993; Vasey, 2006], members of the Mangevo community exhibit a high level of fission–fusion dynamics. In 16 consecutive months, we never found all members of the community together. Rather, community members regularly associated in subgroups that varied in size,

TABLE VI. Results of Linear Mixed Model Predicting Subgroup Cohesion and Using Fruit Availability from Botanical Plots as a Proxy for Fruit Availability. Full Model Results Are Presented Here. The *P* Value Associated With Each Listed Category Indicates Whether It is Significantly Different From the Reference Category Noted in the Footer, e.g., Social is Associated With Lower Group Spread than Feed/Forage

Fixed effect	Estimate	Std. error	<i>t</i> value	<i>P</i>
(Intercept)	1.961	1.116	1.76	0.080
TotalFruitAvailable	−19.180	10.910	−1.76	0.080
ReprodState_Lactation	−1.890	0.718	−2.63	0.009
ReprodState_Nonreproductive	1.573	0.628	2.50	0.013
Activity_Other	0.015	0.235	0.06	0.950
Activity_Rest	−0.581	0.120	−4.86	<0.001
Activity_Social	−2.769	0.607	−4.56	<0.001
Activity_Travel	2.511	0.206	12.18	<0.001
SubGpType_MaleOnly	2.058	0.439	4.69	<0.001
SubGpType_MixedSex	1.382	0.171	8.09	<0.001
SubGpType_SolitaryFemale	5.568	1.149	4.85	<0.001
SubGpType_SolitaryMale	5.596	2.294	2.44	0.015
Subgroup size	2.529	0.085	29.73	<0.001

Model AICc = 96050.5; logLikelihood = −48009.25

The focal individual and data collection day were used as random effects. Gestation is the reference category for the Reproductive State variable.

Feed/forage is the reference category for the Activity variable. Female only is the reference category for the Subgroup Type variable.

composition, and cohesion. On average, subgroup transitions (either fissions or fusions) occurred every 90 min and, much like previous studies of the species [Morland, 1991a,b; Ratsimbazafy, 2002], Mangevo community members spent nearly half of their time alone. When located with other community members, subgroups were generally small (2–7 individuals; 10.6% of community) and cohesive (within 8 m) and typically comprised only one or two subadult and/or adult individuals. Mixed-sex subgroups predominated, followed by male-only and female-only subgroups. Mixed-sex subgroups were also significantly larger and less cohesive than any other subgroup type. That said, with the exception of subadult males, who exhibited unusually strong patterns of association with each other—perhaps representing members of a single birth cohort—adult females appeared to be central to social associations within the community. Females interacted equally with both male and female community members, whereas adult males appeared to interact primarily with adult females. These findings parallel Morland's accounts from Nosy Mangabe [1991a,b], wherein the highest rates of interaction were found among immature age–sex classes and female–female dyads. Affiliative interactions between other age–sex combinations comprised only 15% of the total interactions observed, leading Morland [1991a] to conclude that ruffed lemurs exhibit a type of “female bonded fission–fusion.” Social network analyses among members of the Mangevo ruffed lemur community are underway and will be used to test this hypothesis.

Compared to other primates for which fission–fusion has been quantified, ruffed lemur community size (range: 11–31 individuals) is smaller than that of many chimpanzee communities (e.g., Gombe, Ngogo, Tai), but similar in size to communities observed for bonobos, spider monkeys, and several other platyrrhines (e.g., muriquis, capuchins; Table VII). As is typical of haplorrhine taxa showing high fission–fusion dynamics, ruffed lemur subgroups varied frequently in size, composition, and cohesion. Subgroup duration observed in this study (90 min) was on the same order as in chimpanzees (17–126 min: Boesch [1996]; Boesch & Boesch-Achermann [2000]; Halperin [1979]; Lehmann & Boesch [2004]; Sugiyama [1988]; Reynolds [2005]), bonobos (86–102 min: Kuroda [1979]; White [1988, 1989]), and spider monkeys (120 min: Chapman [1988]), all which are much more fluid in their grouping patterns than species that exhibit low to moderate fission–fusion dynamics, such as hamadryas baboons (e.g., one-male units coalesce just once in 24 hr to sleep [e.g., Swedell & Plummer, 2012]), or snub-nosed monkeys (e.g., groups fission during periods of resource scarcity for periods lasting six or more days [e.g., Ren et al., 2012]).

While these aspects of ruffed lemur fission–fusion resemble patterns found in haplorrhines,

ruffed lemurs also differ quite dramatically in several important respects, particularly in their patterns of subgrouping and social association. For example, despite their overall similarity in community size, both absolute (mean = 2.5 individuals) and relative (10.6%) subgroup sizes at Mangevo were small compared to all other well-studied species, including chimpanzees (range: 4–10 individuals; relative subgroup size mean = 16%), bonobos (4–22 individuals; relative subgroup size mean = 42.9%), and spider monkeys (range: 3–8 individuals; relative subgroup size mean = 14.3%), with the exception of muriquis (range: 2–5 individuals; relative subgroup size mean = 9.9%) (Table VII). Moreover, ruffed lemurs spent more time alone than did any other gregarious primate taxon reported (Table VIII). Whereas most well-studied fission–fusion species typically spend anywhere from 4.2% to 28% of their time alone, ruffed lemurs at Mangevo were without social partners nearly half of the time they were observed (48%), a pattern also described in the Nosy Mangabe community [Morland, 1991a]. The striking amount of time individuals spent solitary led Morland [1991a] to liken ruffed lemur social organization to a combination of the dispersed social systems of the nocturnal cheirogaleids [e.g., reviewed in Müller & Thalmann, 2000] and the fluid fission–fusion groupings of chimpanzees and spider monkeys. Vasey [2006] further refined this assessment by documenting the ranging and spatial association patterns of male and female red ruffed lemurs on Masoala. She found that males resembled non-gregarious nocturnal strepsirrhines in some respects (e.g., minimal overlap between male ranges) whereas females differed substantially from them in that their individual home ranges were larger than those of males and overlapped with those of multiple other females and males within their community. It is somewhat surprising that ruffed lemurs spend such a significant proportion of their time alone, as fission–fusion has long been argued to evolve when the threat of predation is lifted, either by large body size or low density predator communities [Dunbar, 1988; Kummer, 1971; Symington, 1988a; Terborgh & Janson, 1986; Wrangham, 1980]. While they are the largest extant lemurid [Fleagle, 2013], ruffed lemurs are not particularly large [Baden et al., 2008] and the predator community at Ranomafana is not only intact, but diverse [Karpanty & Wright, 2007]. Therefore, the selective pressures leading to this particular social strategy are seemingly incongruent with the hypothesis that fission–fusion results only in the perceived absence of predation threat. Recent work, however, has shown that average predation rates by fossa—among ruffed lemurs' top predators—are low, albeit temporally variable (i.e., predation events are clumped in time) [Irwin et al., 2009]. Thus, it is possible that, while predation can significantly impact ruffed lemur populations [e.g.,

TABLE VII. Cross-Taxonomic Comparisons of Community Size, and Absolute and Relative Subgroup Size

Study site	Community size	Subgroup size		N units	Method	Reference
		Absolute ($\bar{x} \pm SD$)	Relative (%)			
<i>Chimpanzees</i>						
Bossou, Guinea	20	4.00 \pm 2.41	20.0	426 days	1st observation/day	Sakura [1994]
Bossou, Guinea	19–21	5.84 \pm 2.92	29.2	130 days	1st observation/day	Sugiyama [1981]
Budongo Forest, Uganda	46	6.27 \pm 4.42 ^a	13.6	1,824 subgroups	Maximum subgroup size before fission	Newton-Fisher et al. [2000]
Budongo Forest, Uganda	46	5.70 \pm 3.48	12.4	3,164 scans	15 min scans	Newton-Fisher et al. [2000]
Budongo Forest, Uganda	46	5.01 \pm 3.57 ^a	10.9	5,214 scans	30 min scans	Newton-Fisher et al. [2000]
Budongo Forest, Uganda	59	7.01 \pm 4.99	11.9	854 days	Cumulative subgroup size/day (feeding subgroup)	Reynolds [2005]
Budongo Forest, Uganda	59	5.01 \pm 3.57	8.5	5,214 scans	30 min scans	Reynolds [2005]
Fongoli, Senegal	35	15.00 ^a	42.9	unk.	Cumulative subgroup size/day	Pruetz & Bertolani [2009]
Gombe N.P., Tanzania	57	5.60	9.8	498	unk.	Goodall [1986]
Goulougo Triangle Moto, DRC	–	4.85 \pm 4.17 ^b	–	493 days	Cumulative subgroup size/day	Morgan [2007]
Kalinzu Forest—M group, Uganda	55	5.90	13.0	53 days	Cumulative 1 hr subgroup size/day	Mulavwa et al. [2008]
Kibale N.P., Kanyawara, Uganda	–	5.11	–	2,414 subgroups	Count/change in subgroup membership	Chapman et al. [1994, 1995]
Kibale N.P., Kanyawara, Uganda	–	5.24	–	8,190 scans	15 min scans	Chapman et al. [1994, 1995]
Kibale N.P., Kanyawara, Uganda	–	7.13	–	1,208 subgroups	Count/1 hr before and after change in subgroup membership	Chapman et al. [1994, 1995]
Kibale N.P., Kanyawara, Uganda	44	5.60	13.0	1,050 scans	30 min scans	Wrangham et al. [1992, 1996]
Kibale N.P., Ngogo, Uganda	>140	10.27 \pm 10.24	7.4	827 days	1st observation/day	Mitani et al. [2002a]
Kibale N.P., Ngogo, Uganda	145	6.68 \pm 7.02	4.6	3,393 scans	30 min scans	Wakefield [2008, 2013]
Kahuzi–Biega N.P., DRC	22	4.43 \pm 2.50 ^a	20.1	73 subgroups	Count/change in subgroup membership	Basabose [2004]
Mahale Mountain N.P., Tanzania	29	6.20	21.0	218 subgroups	Count/subgroup attending feeding station	Nishida [1968, 1979]
Mahale Mountain N.P., Tanzania	–	12.70 ^b	30.0	unk.	1st observation/day	Matsumoto-Oda et al. [1998]
Mahale Mountain N.P., Tanzania	33.4	15.50 \pm 11.60	13.0	1,167 subgroups	Count/change in subgroup membership	Itoh & Nishida [2007]
Mahale Mountain N.P., Tanzania	33.4	4.00 \pm 3.20	13.0	3,818 scans	10 min scans	Itoh & Nishida [2007]
Rubondo Island, Tanzania	–	3.29 \pm 0.24	–	56 subgroups	Count/change in subgroup membership	Moscovice et al. [2007]
Rubondo Island, Tanzania	–	3.42 \pm 0.29	–	138 subgroups	Count of nests/nest site	Moscovice et al. [2007]
Tai N.P., Côte d'Ivoire	70	3.45 \pm 1.18	4.9	395 scans	1 hr scans	Doran [1997]
Tai N.P., Côte d'Ivoire	76	8.30 ^a	11.0	2,912 subgroups	Count/change in subgroup membership	Boesch [1996]

TABLE VII. Continued

Study site	Community size	Subgroup size		N units	Method	Reference
		Absolute ($\bar{x} \pm SD$)	Relative (%)			
Tai N.P., Côte d'Ivoire	76	10.10 ^a	13.2	527 subgroups	Count/change in subgroup membership	Boesch [1991]; Boesch & Boesch-Achermann [2000]
<i>Bonobos</i>						
Kalinzu Forest—M group, Uganda	17	8.70 \pm 2.80	51.2	465 days	Cumulative 1 hr subgroup size/day	Mulavwa et al. [2008]
Lomako Forest, DRC	—	6.20	—	114 days	1st observation/day	White [1988]
Lomako Forest, DRC	—	5.40 \pm 3.85	—	164 subgroups	Count/change in subgroup membership	White [1988]
Lomako Forest, DRC	10	4.33 \pm 1.75	43.3	87 subgroups	Count/change in subgroup membership	White [1988]
Lomako Forest, DRC	22	7.15 \pm 4.51	32.5	26 subgroups	Count/change in subgroup membership	White [1988]
Lomako Forest, DRC	21	9.69 \pm 5.21	46.1	26 subgroups	Count/change in subgroup membership	White [1988]
Lomako Forest, DRC	—	5.40	—	135	unk.	Malenky & Stiles [1991]
Lomako Forest, DRC	—	5.40	—	164	unk.	Chapman et al. [1994]
Lomako Forest, DRC	36	5.80	22.0	248	unk.	Fruth & Hohmann, pers. comm. (In Boesch 1996)
Lomako Forest, DRC	11–14	4.85 \pm 2.63	27.0	827 counts	Counts	Hohmann & Fruth [2002]
Lukuru, DRC	—	6.4 \pm 4.63	—	121	unk.	Myers Thompson [2002]
Wamba, DRC	23	8.70	51.0	unk.	1 hr scans	Mulavwa et al. [2008]
Wamba, DRC	58	16.9 \pm 11.60 ^b	29.0	147 subgroups	Count/change in subgroup membership	Kuroda [1979]
Wamba, DRC	—	22.70	—	62 counts	Counts	Idani [1991]
<i>Spider monkeys</i>						
Barro Colorado Island, Panama	24	3.44	14.3	unk.	unk.	Milton [1981, 1993]
Manu N.P., Peru	38.5	3.88	10.1	unk.	3 min scans (feeding subgroup)	Symington [1988a,b]
Maracá Ecological Station, Brazil	22	3.01 ^b	13.7	unk.	15 min scans (feeding subgroup)	Nunes [1998]
Punta Laguna, México	28.5	3.85	13.5	unk.	20 min scans	Ramos-Fernandez & Ayala-Orozco [2003]
Tinigua N.P., La Macarena, Colombia	17–18	6.50 \pm 4.10	—	139 subgroups	Count/change in subgroup membership	Shimooka [2003]
Tinigua N.P., La Macarena, Colombia	17–18	5.00 \pm 3.80	—	197 subgroups	Count/change in subgroup membership	Shimooka [2003]
Tinigua N.P., La Macarena, Colombia	17–18	3.70 \pm 2.90	—	164 subgroups	Count/change in subgroup membership	Shimooka [2003]
Tinigua N.P., La Macarena, Colombia	24.5	3.50	10.6	unk.	unk.	Klein & Klein [1977]

TABLE VII. Continued

Study site	Community size	Subgroup size		N units	Method	Reference
		Absolute ($\bar{x} \pm SD$)	Relative (%)			
Noel Kempf Mercado N.P., Bolivia	55	7.80	14.2	unk.	unk.	Wallace [1998]
Noel Kempf Mercado N.P., Bolivia	55	7.52 \pm 6.29	–	–	Count/change in subgroup membership 10 min scans	Wallace [2008]
Santa Rosa N.P., Costa Rica	40	4.94 \pm 4.18	12.4	1,018 scans	Count/change in subgroup membership 10 min scans	Chapman [1990]; Chapman et al. [1995]
Santa Rosa N.P., Costa Rica	26	7.30 \pm 3.50	28.1	137 subgroups	Count/change in subgroup membership	Asensio et al. [2008, 2009]
Yasuni N.P., Ecuador	–	5.50 \pm 2.50	–	5,258	unk.	Suarez [2003]
Yasuni N.P., Ecuador	–	4.90 \pm 3.50	–	9,207 scans	15 min scans	Link & Di Fiore [2013]
Yasuni N.P., Ecuador	25	3.00	12.0	unk.	unk.	Pozo Rivera [2001, 2004]
<i>Muriquis</i>						
Parque Estadual Carlos Botelho, Brazil	54	3.74 \pm 2.32 ^a	6.9	2,440 subgroups	Count/change in subgroup membership	Coles et al. [2012]
Parque Estadual Carlos Botelho, Brazil	54	5.32 \pm 3.27 ^a	9.9	2,440 subgroups	Count/change in subgroup membership	Coles et al. [2012]
Parque Estadual Carlos Botelho, Brazil	23–31	2.60 \pm 1.84 ^b	–	202 subgroups	Count/change in subgroup membership (feeding, fruit)	de Moraes et al. [1998]
Parque Estadual Carlos Botelho, Brazil	23–31	1.83 \pm 1.18 ^b	–	142 subgroups	Count/change in subgroup membership (feeding, leaves)	de Moraes et al. [1998]
Estação Biológica de Caratinga, Brazil	23–31	4.89 \pm 4.79 ^b	–	63 subgroups	Count/change in subgroup membership (feeding, fruit)	de Moraes et al. [1998]
Estação Biológica de Caratinga, Brazil	23–31	2.82 \pm 2.81 ^b	–	61 subgroups	Count/change in subgroup membership (feeding, leaves)	de Moraes et al. [1998]
<i>Capuchins</i>						
Estação Biológica de Caratinga, Brazil	24–28	12.00 \pm 7.78 ^b	–	148 days	Mean subgroup size/15 days (from 5 min scans)	Lynch-Alfaro [2007]
<i>Uakaris</i>						
Lago Preto Conservation Concession, Peru	100+	43.50 \pm 24.10 ^a	–	759 hours	Maximum size/hour	Bowler & Bodmer [2009]
<i>Howler monkeys</i>						
Ometepe Biological Research Station, Nicaragua	–	6.36 \pm 3.27 ^b	–	422 subgroups	Count/change in subgroup membership	Bezanson et al. [2008]
Ometepe Biological Research Station, Nicaragua	–	5.50 \pm 2.90 ^b	–	596 subgroups	Count/change in subgroup membership	Bezanson et al. [2008]
<i>Macaques</i>						

TABLE VII. Continued

Study site	Community size		Subgroup size		N units	Method	Reference
	size	Absolute ($\bar{x} \pm SD$)	Absolute ($\bar{x} \pm SD$)	Relative (%)			
Djurdjura Massif, Algeria	182	15.00–20.00 ^a	–	–	unk. subgroups	Count/change in subgroup membership	Ménard et al. [1990]; Ménard [2002]
<i>Geladas</i>							
Simien Mountains N.P., Ethiopia	–	160.00 ^b	–	–	unk.	1st observation/day	Snyder-Mackler et al. [2012]
<i>Snub-nosed monkeys</i>							
Baimaxueshan National N.R., China	331–480	199.00 ± 63.00 ^b	–	–	24 subgroups	Count/change in subgroup membership	Ren et al. [2012]
<i>Ruffed lemurs</i>							
Nosy Mangabe S.R., Madagascar	11–16	2.50	–	–	84 days	1st observation/day	Morland [1991a]
Nosy Mangabe S.R., Madagascar	11–16	2.80 ± 0.26	–	–	60 subgroups	Subgroups encountered during census	Morland [1991a]
Masoala N.P., Andranobe, Madagascar	18–31	2.60 ± 0.55	–	–	5 subgroups	Number of 'core groups'	Vasey [2006]
Ranomafana N.P., Mangevo, Madagascar	24	2.54 ± 0.85	10.6	–	40,840 scans	5 min scans	This study

Note: Subgroup size includes only independent (adult and subadult) members unless otherwise noted.
^aStudies where dependent offspring (juveniles or infants) are included in estimates of subgroup size.

four predation events in three days within the same ruffed lemur community; Baden, unpublished data], the perceived threat is infrequent enough that selection for greater group cohesion is weak.

Beyond general differences associated with subgroup type, ruffed lemurs also differ in their preferences for social associates. In both chimpanzee and spider monkey societies, males are typically thought of as being at the core of social dynamics [e.g., Gilby & Wrangham, 2008; Machanda et al., 2013; Slater et al., 2009; Symington, 1987; Wrangham & Smuts, 1980]. Male–male affiliative bonds tend to be stronger than either mixed-sex or female–female bonds [Fedigan & Baxter, 1984; Gilby & Wrangham, 2008; Machanda et al., 2013; Shimooka, 2003; Slater et al., 2009; but see Lehmann & Boesch, 2008; Williams et al., 2002]. Strong affiliative interactions between males have been said to play an important role in maintaining the cooperation needed in a context of high intergroup competition [Gilby & Wrangham, 2008; Link et al., 2009], as it potentially provides benefits to the resident males associated with defense and access to resources and mates [van Hooff & van Schaik, 1994], and in cooperatively competing with rival neighboring groups [Link, 2011]. While it is difficult to compare patterns of association strength across studies (due to incongruous methods of quantifying affiliation), it is clear that, contrary to the male-biased relationships observed in chimpanzees and spider monkeys, patterns of ruffed lemur association are predominantly female-centered [Morland, 1991a; this study (Fig. 4)], mirroring instead those patterns observed in bonobos [e.g., Hohmann & Fruth, 2002]. Following the same reasoning as described above for chimpanzees and spider monkeys, strong female bonds may help to facilitate cooperation during territory defense [Baden, unpublished data; Morland, 1991a; Vasey, 2006]. However, unlike chimpanzees and spider monkeys, which are male-philopatric (and, therefore, are more likely to be genetic relatives, but see [Vigilant et al., 2001; Di Fiore, 2009; Di Fiore et al., 2009]), the strong female-biased relationships observed in ruffed lemurs [Morland, 1991a; this study] are unlikely to be driven by kinship, as both males and females disperse [Baden, 2011; Baden et al., 2014; Razakamaharavo et al., 2010]. Therefore, it is unclear as to why female associations are stronger than those among males. One explanation may be in their communal breeding reproductive strategy. Because infants cannot cling at birth, litters are parked in nests and tree tangles until capable of independent travel, and it is during this time that evidence of communal breeding has been reported, including use of communal nests (crèches) and cooperative infant care (e.g., babysitting, allomaternal nursing) [Baden, 2011; Baden et al., 2013; Morland, 1989, 1990, 1991a; Tecot et al., 2013; Vasey, 2007]. Although infant care is not limited to

TABLE VIII. Cross-Taxonomic Comparison of Subgroup Type (% of Observations)

Site	Solitary				Mixed	Reference
	Overall	Male	Female	Male		
<i>Chimpanzees</i>						
Assirik	19.0	–	–	2.0	18.0	Tutinet al. [1983]
Bossou, Guinea	–	–	–	–	49.0	Sakura [1994]
Budongo Forest, Uganda	–	–	–	8.7	3.1	Reynolds [2005]
Budongo Forest, Uganda (travel)	–	27.9	15.0	26.4	3.6	Sugiyama [1968]
Budongo Forest, Uganda (feed)	–	12.8	13.7	24.8	4.7	Sugiyama [1968]
Gombe N.P., Tanzania	18.0	–	–	10.0	24.0	Goodall [1968]
Kibale N.P., Kanyawara, Uganda	28.0	–	–	–	–	Chapman et al. [1995]
Kibale N.P., Kanyawara, Uganda	–	3.8	19.1	16.6	40.4	Chapman et al. [1995]
Kibale N.P., Ngogo, Uganda	–	–	–	–	19.7	[2008, 2013]
Mahale Mountain N.P., Tanzania	–	9.0	2.0	9.0	17.0	Matsumoto-Oda et al. [1998]
Mahale Mountain N.P., Tanzania	21.0	–	–	11.0	13.0	Nishida [1968]
Tai N.P., Côte d'Ivoire	10.0	6.0	4.0	20.0	18.0	Boesch [1996]
<i>Bonobos</i>						
Lomako Forest, DRC	14.0	–	–	5.0	5.0	White [1988, 1992]
Lomako Forest, DRC	–	2.3	2.1	8.0	21.0	Hohmann & Fruth [2002]
Wamba, DRC	6.1	–	–	2.5	4.9	Kuroda [1979]
<i>Spider monkeys</i>						
Santa Rosa N.P., Costa Rica	–	8.0	24.0	3.0	47.0	Chapman [1990]
<i>Capuchins</i>						
Estação Biológica de Caratinga, Brazil	–	2.7	3.4	6.8	1.4	Lynch-Alfaro [2007]
<i>Howler monkeys</i>						
Ometepe Biological Research Station, Nicaragua	4.2	–	–	0.0	0.0	Bezanson et al. [2008]
<i>Muriquis</i>						
Parque Estadual Carlos Botelho, Brazil	–	2.0	5.0	13.4	1.2	Coles et al. [2012]
<i>Ruffed lemurs</i>						
Nosy Mangabe S.R., Madagascar	46.0	16.2	18.2	5.4	7.4	Morland [1991a]
Ranomafana N.P., Mangevo, Madagascar	46.0	20.7	23.5	6.8	4.7	This study

Every attempt was made to standardize definitions of subgroup types across studies. However, some studies used subgroup types that were not obviously equivalent to our own (e.g., “adult subgroup”) and were, thus, excluded here. In these cases, rows will not sum to 100%. Solitary female category includes mothers with dependent young.

mothers [Baden et al., 2013; Morland, 1990; Vasey, 2007], there is a strong bias toward reproductive females [Baden, unpublished data], and thus maintaining strong social relationships year-round may facilitate cooperative care during the time of year when infants are dependent.

Ruffed lemurs are notable among fission–fusion taxa in that their fluid grouping patterns display both daily and seasonal components. Indeed, not only do ruffed lemur subgroups vary in size, composition, and cohesion throughout the course of a day—patterns that we have demonstrated largely resemble those of haplorrhine taxa (Tables VII and VIII)—our multivariate analyses also revealed significant, combined effects of climate, fruit availability, and female reproductive state on ruffed lemur fission–fusion dynamics. For example, we found that ruffed lemur subgroups were larger and more cohesive during the warm-wet season (Feb–May), a time that largely coincides with high fruit availability (Fig. 1). As resources became increasingly scarce—moving into the cool-wet months of June, July, and August—subgroups in our study became smaller and less cohesive. Subgroups were also smaller at lower temperatures and during periods of reduced rainfall, variables that have been closely linked with reductions in plant productivity [Dewar & Richard, 2007; Hemingway, 1996, 1998; Meyers & Wright, 1993; Overdorff, 1993]. These results are consistent with those of Morland [1991a,b], who described ruffed lemurs as having fewer inter-individual associations and spending more time alone during cool periods of resource scarcity, and with Vasey [2006], who documented the dispersion of core groups into separate core areas during both gestation and the food-scarce, cold rainy season. Together, these observations support the hypothesis that fission–fusion dynamics primarily function to reduce feeding competition in the face of fruit scarcity [Anderson et al., 2002; Chapman et al., 1995; Lehmann et al., 2007; Shimooka, 2003].

Ruffed lemur fission–fusion dynamics also varied with reproductive season. In particular, subgroups were smallest and most cohesive during the months associated with lactation and high infant dependence (October–December). This pattern was, in large part, probably driven by the initial six weeks post-partum, which comprised the period of natal nest use and exclusive maternal care [Baden, 2011; Baden et al., 2013]. Similarly, Morland [1991a] and Vasey [2006] have both noted that reproductive state influenced female gregariousness, with females in early lactation being more solitary than non-lactating females in the same months. While these patterns are not unlike observations in chimpanzees, whereby reproductive state determines much of a female's sociality [Goodall, 1986; Mitani et al., 2002b; Wakefield, 2013], they differ in that patterns of ruffed lemur fission–fusion take on both a daily and

seasonal form. Like most strepsirrhines, ruffed lemurs stand apart from chimpanzees and all other haplorrhines in their patterns of strict seasonal breeding and reproductive synchrony [Baden et al., 2013; Bogart et al., 1977a,b; Boskoff, 1977a; Rasmussen, 1985; Morland, 1993]. All females within a community generally come into vaginal estrus once during a brief, synchronized 24–72 hour period each year (i.e., the “reproductive season”) [Foerg, 1982; Baden, unpublished data]. During all other times of year, female genitalia are imperforate [Foerg, 1982]. Because reproduction in this taxon is so tightly synchronized among individuals, it therefore results in a corresponding behavioral synchrony across seasons (e.g., reduced fission–fusion dynamics during early lactation). These patterns support claims that ruffed lemur social organization appears to be largely driven by the unique life history of this genus [Vasey, 2003, 2007]. We should note that the relationships described herein for the Mangevo population are derived from the first three months of lactation only. A highly gregarious period during the latter half of lactation has been described elsewhere [Vasey, 2007].

To our knowledge, these seasonal patterns of fission–fusion dynamics are unique to ruffed lemurs and appear to be consistent across sites and study periods (despite spanning several decades). It is worth noting that the few studies where ruffed lemur social dynamics do not conform to these patterns have been either very short in duration (e.g., two months during Madagascar's austral winter when subgroups are smaller and more cohesive: White [1991]), or of populations that have faced recent ecological hardships (e.g., Cyclone Gretelle significantly impacted forest structure and fruit availability: Ratsimbazafy [2002]; long-term selective logging removed large fruiting trees and affected average tree height, diameter, and density: Balko [1998]; Balko and Underwood [2005]). Thus, despite their obligate frugivory and sensitivity to habitat disturbances, ruffed lemur social flexibility has likely allowed them to persist under suboptimal ecological conditions, at least in the short term.

While the most common assumption is that fission–fusion parameters strongly depend on ecological variables and reproductive strategies of the species, demographic variables (e.g., population density, community size, number of males) can also affect subgrouping patterns [Boesch, 1996; Lehmann & Boesch, 2004]. For example, several studies have found an inverse relationship between community size and relative party size; at smaller community size, sexes showed greater cohesiveness and reduced fluidity (lower transition rates) [Boesch & Boesch-Achermann, 2000; Lehmann & Boesch, 2004; Sugiyama & Koman, 1979]. At present, the data available from ruffed lemurs appear to also fit this pattern (Table IX). At sites where ruffed lemur

TABLE IX. Comparison of Ruffed Lemur Community Size, Sex Ratio, and Absolute and Relative Subgroup Sizes Across Sites and Long-Term Studies

Species	Study site	Population density/km ²	Social organization	Community size	Sex ratio	Subgroup size			Reference
						Absolute ($\bar{x} \pm SD$)	Relative (%)	Study duration (months)	
<i>Varecia rubra</i>	Ambatonakolahy (Masoala N.P.)	21.5–23.3	Fission–fusion	5.5 (n = 2)	1.50	unk.	unk.	11	Rigamonti [1993]
<i>Varecia rubra</i>	Andranobe (Masoala N.P.) ^a	31.2–53.4	Fission–fusion	13.0 (n = 1)	0.38	2.60 ± 0.55	23.64	12	Vasey [1997, 2006]
<i>Varecia variegata</i>	Nosy Mangabe	29.0–43.0	Fission–fusion	11.0 (n = 2)	1.02	2.80 ± 0.26	21.54	13	Morland [1991a,b]
<i>Varecia variegata</i>	Vatoharanana (Ranomafana N.P.)	1.5	Dispersed monogamy	2.0 (n = 1)	1.00	2.00	100	2	White [1989, 1991]
<i>Varecia variegata</i>	Vatoharanana (Ranomafana N.P.)	3.0–6.0	Cohesive mm:mf	5.7 (n = 3)	1.44	5.67 ± 1.15	100	18	Balko [1998]
<i>Varecia variegata</i>	Mangevo (Ranomafana N.P.)	23.4	Fission–fusion	24.0 (n = 1)	1.38	2.54 ± 0.85	10.59	16	Baden [2011], this study
<i>Varecia variegata</i>	Manombo (Ranomafana N.P.)	0.25	Dispersed mm:mf	2.4 (n = 2)	0.83	2.42 ± 0.59	100	18	Ratsimbazafy [2002]

Community and subgroup sizes include adults and subadults only (infants excluded). For studies where fission–fusion was absent, community size is equivalent to subgroup size.

Sex ratios are presented as the number of adult males to every one adult female.

^aAbsolute and relative subgroup sizes based on core group sizes.

population densities were low (0.25–6 ind./km²), animals reportedly associated in small, stable pair-bonded or multimale-multifemale groups (but see discussion above regarding limitations to these studies). By contrast, animals living at higher population densities (>20 ind./km²) tended to associate in larger, more fluid multimale-multifemale groups that were characterized by both daily and seasonal patterns of fission–fusion social dynamics. These latter communities were all similar in size and sex ratio, and subgroup dynamics (e.g., average absolute and relative subgroup sizes) were remarkably similar. Unfortunately, further comparisons were not possible because comparable data were not available from all sites. In the future, it will be informative to compare the fission–fusion dynamics (e.g., transition rates, patterns of group size, and cohesion) in ruffed lemurs at sites with different population densities and among communities differing in size.

In summary, we found that black-and-white ruffed lemurs display a distinct pattern of fission–fusion dynamics that is both markedly different from and strikingly similar to haplorrhine taxa with fluid fission–fusion societies. Compared to haplorrhines, ruffed lemurs exhibited smaller subgroups, dramatically lower rates of association, and a female-centered social organization. What is more, group dynamics varied seasonally with changes in climate, fruit availability, and reproductive state. These seasonal patterns of ruffed lemur fission–fusion paired with aspects of their spatial ecology (e.g., shorter travel distances, smaller ranges, and fewer associations during the austral winter: [Baden & Gerber, in prep; Vasey, 2006]) and unique life histories (e.g., litters, seasonal breeding, and boom-bust reproduction: [Baden et al., 2013; Vasey, 2007]) are consistent with Wright’s [1999] energy frugality hypothesis, which postulates that lemur traits are adaptations to conserve energy and/or to maximize use of scarce resources. However, while fission–fusion dynamics may comprise one strategy among a suite of characteristics making ruffed lemurs uniquely adapted to their harsh and unpredictable island environment, similarities between ruffed lemurs and other fluid fission–fusion species suggest that the factors shaping these patterns can select for remarkably similar behavioral strategies, particularly in convergent settings.

ACKNOWLEDGEMENTS

We would like to thank Anthony Di Fiore, Natalie Vasey, and one anonymous reviewer for their thorough and insightful comments and critiques on earlier versions of this manuscript. Thanks to ICTE, MICET and Centre ValBio for facilitating research in Madagascar, and ANGAP/Madagascar’s National Parks for permission to conduct research in

RNP. Drs. Randy Junge, Felicia Knightly, Angie Simai, Edward E. Louis, Jr., and staff of the Madagascar Biodiversity Project provided invaluable assistance during capture procedures. Reycheil Chadwick, Lindsay Dytham, A.J. Lowin, Solo Justin, Telo Albert, Lahitsara Pierre, Razafindrakoto Georges, Leroa, and Velomaro provided field assistance, companionship, and advice on the ground. Funding was generously provided by National Science Foundation DDIG (BSC-0725975), US Department of State Fulbright Fellowship, The Leakey Foundation, Primate Conservation, Inc., Conservation International's Primate Action Fund, Stony Brook University, and Henry Doorly Zoo. Research complied with the laws and guidelines set forth by ANGAP/Madagascar National Parks and Stony Brook University's IACUC (#2005-20081449).

REFERENCES

- Alexander RD. 1974. The evolution of social behavior. *Annual Review of Ecological Systems* 5:325–383.
- Altmann J. 1974. Observational study of behavior: sampling methods. *Behaviour* 49:227–267.
- Anderson DP, Nordheim EV, Boesch C, Moermond TC. 2002. Factors influencing fission–fusion grouping in chimpanzees in the Tai National Park, Cote d'Ivoire. In: Boesch C, Hohmann G, Marchant L, editors. *Behavioral diversity in chimpanzees and bonobos*. Cambridge: Cambridge University Press. p 90–101.
- Asensio N, Korstjens AH, Schaffner CM, Aureli F. 2008. Intragroup aggression, fission–fusion dynamics and feeding competition in spider monkeys. *Behaviour* 145:983–1001.
- Asensio N, Korstjens AH, Schaffner CM, Aureli F. 2009. Intragroup aggression, fission–fusion dynamics and feeding competition in spider monkeys. *Behaviour* 145:983–1001.
- Aureli F, Schaffner CM. 2008. Social interactions, social relationships and the social system of spider monkeys. In: Campbell, editor. *Spider monkeys: behavior, ecology and evolution of the genus Ateles*. Cambridge: Cambridge University Press. p 236–265.
- Aureli F, Schaffner CM, Boesch C, et al. 2008. Fission–fusion dynamics. *Current Anthropology* 49:627–654.
- Baden AL. 2011. *Communal infant care in black-and-white ruffed lemurs (Varecia variegata)* [dissertation]. Stony Brook: Stony Brook University.
- Baden AL, Brenneman RA, Louis EE, Jr. 2008. Morphometrics of wild black-and-white ruffed lemurs [*Varecia variegata*, Kerr 1972]. *American Journal of Primatology* 70:913–926.
- Baden A, Wright P, Louis E, Bradley BJ. 2013. Cooperation confers maternal success in a social primate. *Behavioral Ecology and Sociobiology* 67:1939–1950.
- Baden AL, Holmes SM, Johnson SE, et al. 2014. Species-level view of population structure and gene flow for a critically endangered primate (*Varecia variegata*). *Ecology and Evolution* 4:2675–2692.
- Balko EA. 1998. A behaviorally plastic response to forest composition and logging disturbance by *Varecia variegata variegata* in Ranomafana National Park, Madagascar [dissertation]. State University of New York – College of Environmental Science and Forestry.
- Balko EA, Underwood HB. 2005. Effects of forest structure and composition on food availability for *Varecia variegata* at Ranomafana National Park, Madagascar. *American Journal of Primatology* 66:45–70.
- Bartoń K. 2014. MuMIn: multi-model inference. Model selection and model averaging based on information criteria (AICc and alike). R package version 1.10.5, <http://cran.r-project.org/web/packages/MuMIn/index.html>.
- Basabose AK. 2004. Fruit availability and chimpanzee party size at Kahuzi montane forest, Democratic Republic of Congo. *Primates* 45:211–219.
- Bates D, Maechler M, Bolker B, Walker S. 2014. lme4: linear mixed-effects models using Eigen and S4. R package version 1.1-7, <http://CRAN.R-project.org/package=lme4>.
- Bezanson M, Garber PA, Murphy JT, Premo LS. 2008. Patterns of subgrouping and spatial affiliation in a community of mantled howling monkeys (*Alouatta palliata*). *American Journal of Primatology* 70:282–293.
- Boesch C. 1991. The effects of leopard predation on grouping patterns in forest chimpanzees. *Behaviour* 117:220–242.
- Boesch C. 1996. Social grouping in Tai chimpanzees. In: McGrew W, Merchant L, Nishida T, editors. *Great ape societies*. Cambridge: Cambridge University Press. p 101–113.
- Boesch C, Boesch H. 1989. Hunting behavior of wild chimpanzees in the Tai National Park. *American Journal of Physical Anthropology* 78:547–573.
- Boesch C, Boesch-Achermann H. 2000. *The chimpanzees of the Tai forest: behavioral ecology and evolution*. Oxford: Oxford University Press.
- Bogart MH, Cooper RW, Benirschke K. 1977a. Reproductive studies of black and ruffed lemurs. *International Zoological Yearbook* 17:177–182.
- Bogart MH, Kumamoto AT, Lasley BL. 1977b. A comparison of the reproductive cycle of three species of Lemur. *Folia Primatologica* 28:134–143.
- Boskoff KJ. 1977. Aspects of reproduction in ruffed lemurs (*Lemur variegatus*). *Folia Primatologica* 28:241–250.
- Bowler M, Bodmer R. 2009. Social behavior in fission–fusion groups of red uakari monkeys (*Cacajao calvus ucayalii*). *American Journal of Primatology* 71:976–987.
- Bowler M, Knogge C, Heymann EW, Zinner D. 2012. Multilevel societies in new world primates? Flexibility may characterize the organization of Peruvian Red Uakaris (*Cacajao calvus ucayalii*). *International Journal of Primatology* 33:1110–1124.
- Britt A. 1997. *Environmental influences on the behavioural ecology of the black-and-white ruffed lemur (Varecia variegata variegata, Kerr 1972)* [dissertation]. University of Liverpool.
- Brockman DK, Willis MS, Karesh WB. 1987. Management and husbandry of ruffed lemurs, *Varecia variegata*, at the San Diego Zoo. II. Reproduction, pregnancy, parturition, litter size, infant care, and reintroduction of hand-raised infants. *Zoo Biology* 6:349–363.
- Burnham KP, Anderson D. 2002. *Model selection and multi-model inference*. New York: Springer.
- Caillaud D, Craft MA, Meyers LA. 2013. Epidemiological effects of group size variation in social species. *Journal of the Royal Society Interface* 10:20130206.
- Cairns SJ, Schwager SJ. 1987. A comparison of association indices. *Animal Behavior* 35:1454–1469.
- Chapman CA. 1988. Patch use and patch depletion by the spider and howling monkeys of Santa Rosa National Park, Costa Rica. *Behaviour* 105:99–116.
- Chapman CA. 1990. Association patterns of spider monkeys: the influence of ecology and sex on social organization. *Behavioral Ecology and Sociobiology* 26:409–414.
- Chapman CA. 1990b. Ecological constraints on group size in three species of Neotropical primates. *Folia Primatologica* 55:1–9.
- Chapman CA, Wrangham RW. 1993. Range use of the forest chimpanzees of Kibale: implications for the evolution of chimpanzee social organization. *American Journal of Primatology* 31:263–273.
- Chapman CA, White FJ, Wrangham RW. 1994. Party size in chimpanzees and bonobos. In: Wrangham RW, McGrew WC,

- de Waal FDM, Heltne PG, editors. Chimpanzee cultures. Cambridge, Massachusetts: Harvard University Press.
- Chapman C, Chapman L, Wrangham R. 1995. Ecological constraints on group size: an analysis of spider monkey and chimpanzee subgroups. *Behavioral Ecology and Sociobiology* 36:59–70.
- Clutton-Brock TH, Harvey PH. 1977. Primate ecology and social organization. *Journal of Zoology* 183:1–39.
- Coles RC, Lee PC, Talebi M. 2012. Fission–fusion dynamics in southern muriquis (*Brachyteles arachnoides*) in continuous Brazilian Atlantic Forest. *International Journal of Primatology* 33:93–114.
- Crockett C, Janson CH. 2000. Infanticide in red howlers: female group size, male membership, and a possible link to folivory. In: van Schaik CP, Janson CH, editors. Infanticide by males and its implications. Cambridge: Cambridge University Press. p 75–98.
- Dewar RE, Richard AF. 2007. Evolution in the hypervariable environment of Madagascar. *Proceedings of the National Academy of Sciences* 104:13723–13727.
- Di Fiore A. 2009. Genetic approaches to the study of dispersal and kinship in New World primates. In: Garber PA, Estrada A, Bicca-Marques JC, Heymann EW, Strier KB, editors. South American primates: comparative perspectives in the study of behavior, ecology and conservation. New York: Springer. p 211–250.
- Di Fiore A, Rendall D. 1994. Evolution of social organization: a reappraisal for primates by using phylogenetic methods. *Proceedings of the National Academy of Sciences USA* 91:9941–9945.
- Di Fiore A, Link AL, Schmitt CA, Spehar SN. 2009. Dispersal patterns in sympatric woolly and spider monkeys: integrating molecular and observational data. *Behaviour* 146:437–470.
- Di Fiore A, Link A, Campbell CJ. 2011. The Atelines: behavior and socioecological diversity in a new world monkey radiation. In: Campbell CJ, Fuentes AF, MacKinnon KC, Stumpf R, Bearder S, editors. Primates in perspective. Oxford: Oxford University Press. p 155–188.
- Doran D. 1997. Influence of seasonality on activity patterns, feeding behavior, ranging, and grouping patterns in Tai chimpanzees. *International Journal of Primatology* 18:183–206.
- Dunbar R. 1988. Primate social systems. New York: Cornell University Press.
- Fedigan LM, Baxter MJ. 1984. Sex differences and social organization in free-ranging spider monkeys (*Ateles geoffroyi*). *Primates* 25:279–294.
- Fleagle JG. 2013. Primate adaptation and evolution. San Diego, CA: Elsevier. p 441.
- Foerg R. 1982. Reproductive behavior in *Varecia variegata*. *Folia Primatologica* 38:108–121.
- Furuichi T. 2009. Factors underlying party size differences between chimpanzees and bonobos: a review and hypotheses for future study. *Primates* 50:197–209.
- Garamszegi LZ. 2011. Information-theoretic approaches to statistical analysis in behavioural ecology: an introduction. *Behavioral Ecology and Sociobiology* 65:1–11.
- Ghiglieri MP. 1984. Feeding ecology and sociality of chimpanzees in Kibale Forest, Uganda. In: Rodman PS, Cant JGH, editors. Adaptations for foraging in nonhuman primates. Contributions to an organismal biology of prosimians, monkeys, and apes. New York: Columbia University Press. p 161–194.
- Gilby IC, Wrangham RW. 2008. Association patterns among wild chimpanzees (*Pan troglodytes schweinfurthii*) reflect sex differences in cooperation. *Behavioral Ecology and Sociobiology* 62:1831–1842.
- Ginsberg JR, Young TP. 1992. Measuring association between individuals or groups in behavioural studies. *Animal Behaviour* 44:377–379.
- Glander K. 1993. Capture and marking techniques for arboreal primates. In: Estrada AR-LE, Lopez-Wilchis R, Coates-Estrada R, editors. Estudios Primatológicos En Mexico. Veracruz: Universidad Veracruzana. p 229–304.
- Goodall J. 1968. The behaviour of free-living chimpanzees in the Gombe Stream area. *Animal Behavior Monographs* 1:161–311.
- Goodall J. 1986. The chimpanzees of Gombe. Cambridge: Belknap Press.
- Halperin SD. 1979. Temporary association patterns in free-ranging chimpanzees: An assessment of individual grouping preferences. In: Hamburg DA, McCown ER, editor. The Great Apes. Menlo Park: Benjamin/Cummings.
- Hasegawa T. 1990. Sex differences in ranging patterns. In: Nishida T, editor. The chimpanzees of the Mahale mountains: sexual and life history strategies. Tokyo: University of Tokyo Press. p 99–114.
- Hashimoto C, Furuichi T, Tashiro Y. 2001. What factors affect the size of chimpanzee parties in the Kalinzu Forest, Uganda? Examination of fruit abundance and number of Estrous females. *International Journal of Primatology* 22:947–959.
- Hashimoto C, Suzuki S, Takenoshita Y, et al. 2003. How fruit abundance affects the chimpanzee party size: a comparison between four study sites. *Primates* 44:77–81.
- Hemingway CA. 1996. Morphology and phenology of seeds and whole fruit eaten by Milne-Edwards' sifaka, *Propithecus diadema edwardsi*, in Ranomafana National Park, Madagascar. *International Journal of Primatology* 17:637–659.
- Hemingway CA. 1998. Selectivity and variability in the diet of Milne-Edwards' sifakas (*Propithecus diadema edwardsi*): implications for folivory and seed-eating. *International Journal of Primatology* 19:355–377.
- Hill RA, Lee PC. 1998. Predation risk as an influence on group size in cercopithecoïd primates: implications for social structure. *Journal of the Zoological Society of London* 245:447–456.
- Hohmann G, Fruth B. 2002. Dynamics in social organization of bonobos (*Pan paniscus*). In: Boesch C, Hohmann G, Marchant LF, editors. Behavioural diversity in chimpanzees and bonobos. Cambridge: Cambridge University Press. p 138–150.
- Idani G. 1991. Social relationships between immigrant and resident bonobo (*Pan paniscus*) females at Wamba. *Folia Primatologica* 57:83–95.
- Irwin MT, Raharison JL, Wright PC. 2009. Spatial and temporal variability in predation on rainforest primates: do forest fragmentation and predation act synergistically? *Animal Conservation* 12:220–230.
- Isabirye-Basuta G. 1988. Food competition among individuals in a free-ranging chimpanzee community in Kibale Forest, Uganda. *Behaviour* 105:135–147.
- Isbell LA. 1991. Contest and scramble competition: patterns of female aggression and ranging behavior among primates. *Behavioral Ecology and Sociobiology* 2:143–155.
- Itoh N, Nishida T. 2007. Chimpanzee grouping patterns and food availability in Mahale Mountains National Park, Tanzania. *Primates* 48:87–96.
- Janson CH, Goldsmith ML. 1995. Predicting group size in primates: foraging costs and predation risks. *Behavioral Ecology* 6:326–336.
- Janson CH, van Schaik CP. 1988. Recognizing the many faces of primate food competition: methods. *Behaviour* 105:165–186.
- Kappeler PM. 1997. Determinants of primate social organization: comparative evidence and new insights from Malagasy lemurs. *Biological Reviews* 72:111–151.
- Karpanty SM, Wright PC. 2007. Predation on lemurs in the rainforest of Madagascar by multiple predator species: observations and experiments. In: Gursky-Doyen S, Nekaris KAI, editors. Primate anti-predator strategies. New York, NY: Springer. p 77–99.

- Kawanaka K. 1984. Association, ranging, and the social unit in chimpanzees of the Mahali Mountains, Tanzania. *International Journal of Primatology* 5:411–434.
- Kinzey WG, Cunningham EP. 1994. Variability in platyrrhine social organization. *American Journal of Primatology* 34:185–198.
- Klein LL, Klein DB. 1977. Feeding behaviour of the Colombian spider monkey. In: Clutton-Brock TH, editor. *Studies of feeding and ranging behaviour in lemurs, monkeys and apes*. New York, NY: Academic Press. p 153–181.
- Koenig A. 2002. Competition for resources and its behavioural consequences among female primates. *International Journal of Primatology* 23:759–783.
- Kummer H. 1971. Primate societies: group techniques of ecological adaptation. Chicago: Aldine. p 160.
- Kuroda S. 1979. Grouping of the pygmy chimpanzee. *Primates* 20:161–183.
- Kuznetsova A, Brockhoff PB, Christensen RHB. 2014. lmerTest: tests for random and fixed effects for linear mixed effect models. R package version 2.0-11, <http://cran.r-project.org/web/packages/lmerTest/index.html>.
- Lehmann J, Boesch C. 2004. To fission or to fusion: effects of community size on wild chimpanzees (*Pan troglodytes verus*) social organization. *Behavioral Ecology and Sociobiology* 56:207–216.
- Lehmann J, Boesch C. 2008. Sexual differences in chimpanzee sociality. *International Journal of Primatology* 29:65–81.
- Lehmann J, Korstjens AH, Dunbar RIM. 2007. Fission–fusion social systems as a strategy for coping with ecological constraints: a primate case. *Evolutionary Ecology* 21:613–634.
- Link A. 2011. Social and ecological determinants of “fission fusion” sociality and grouping strategies in the white bellied spider monkey (*Ateles belzebuth belzebuth*) in a lowland rainforest in Western Amazonia [dissertation]. New York: New York University.
- Link A, Di Fiore A, Spehar SN. 2009. Female-directed aggression and social control in spider monkeys. In: Muller MN, Wrangham RW, editors. *Sexual coercion in primates and humans: an evolutionary perspective on male aggression against females*. Cambridge: Harvard University Press. p 157–183.
- Lynch-Alfaro JW. 2007. Subgrouping patterns in a group of wild *Cebus apella nigrinus*. *International Journal of Primatology* 28:271–289.
- Machanda ZP, Gilby IC, Wrangham RW. 2013. Male-female association patterns among free-ranging chimpanzees (*Pan troglodytes schweinfurthii*). *International Journal of Primatology* 34:917–938.
- Majolo B, Vizioli ADB, Schino G. 2008. Costs and benefits of group living in primates: group size effects on behaviour and demography. *Animal Behaviour* 76:1235–1247.
- Malenky RK, Stiles EW. 1991. Distribution of terrestrial herbaceous vegetation and its consumption by *Pan paniscus* in the Lomako Forest, Zaire. *American Journal of Primatology* 23:153–169.
- Mantel N. 1967. The detection of disease clustering and a generalized regression approach. *Cancer Research* 27:209–220.
- Marlowe FW. 2005. Hunter-gatherers and human evolution. *Evolutionary Anthropology: Issues, News, and Reviews* 14:54–67.
- Matsumoto-Oda A. 1999. Mahale chimpanzees: grouping patterns and cycling females. *American Journal of Primatology* 47:197–207.
- Matsumoto-Oda A, Hosaka K, Huffman MA, Kawanaka K. 1998. Factors affecting party size in chimpanzees of the mahale mountains. *International Journal of Primatology* 19:999–1011.
- Ménard N. 2002. Ecological plasticity of barbary macaques (*Macaca sylvanus*). *Evolutionary Anthropology* 1:95–100.
- Ménard N, Hecham R, Vallet D, Chickhi H, Gautier-Hion A. 1990. Grouping patterns of a mountain population of *Macaca sylvanus* in Algeria – a fission–fusion system? *Folia Primatologica* 55:166–175.
- Meyers DM, Wright PC. 1993. Resource tracking: food availability and *Propithecus* seasonal reproduction. In: Ganzhorn J, Kappeler PM, editors. *Lemur social systems and their ecological basis*. New York: Plenum Press. p 179–192.
- Milton K. 1981. Food choice and digestive strategies of two sym-patric primate species. *American Naturalist* 117:496–505.
- Milton K. 1993. Diet and social organization of a free-ranging spider monkey population: the development of species typical behaviour in absence of adults. In: Pereira M, Fairbanks LA, editors. *Juvenile primates*. Oxford: Oxford University Press. p 173–181.
- Mitani JC, Watts DP, Lwanga JS, Marchant L. 2002a. Ecological and social correlates of chimpanzee party size and composition. In: Boesch C, Hohmann G, Boesch C, Hohmann G, editors. *Behavioural diversity in chimpanzees and bonobos*. Cambridge: Cambridge University Press. p 102–111.
- Mitani JC, Watts DP, Muller MM. 2002b. Recent developments in the study of wild chimpanzee behavior. *Evolutionary Anthropology* 11:9–25.
- Morgan D. 2007. Socio-ecology of chimpanzees in the Goulougo Triangle, Republic of Congo [dissertation]. Cambridge: University of Cambridge.
- Morland HS. 1989. Infant survival and parental care in ruffed lemurs (*Varecia variegata*) in the wild. *American Journal of Primatology* 18:157.
- Morland HS. 1990. Parental behavior and infant development in ruffed lemurs (*Varecia variegata*) in a Northeast Madagascar rain forest. *American Journal of Primatology* 20:253–265.
- Morland H. 1991a. Social organization and ecology of black and white ruffed lemurs (*Varecia variegata variegata*) in lowland rain forest, Nosy Mangabe, Madagascar [dissertation]. New Haven: Yale University.
- Morland HS. 1991b. Preliminary report on the social organization of ruffed lemurs (*Varecia variegata variegata*) in a northeast Madagascar rain forest. *Folia Primatologica* 56:157–161.
- Morland HS. 1993. Reproductive activity of ruffed lemurs (*Varecia variegata variegata*) in a Madagascar rain forest. *American Journal of Physical Anthropology* 91:71–82.
- Moscovice LR, Issa MH, Petrzelkova KJ, et al. 2007. Fruit availability, chimpanzee diet, and grouping patterns on Rubondo Island, Tanzania. *American Journal of Primatology* 69:487–502.
- Müller AE, Thalmann U. 2000. Origin and evolution of primate social organisation: a reconstruction. *Biological Review of the Cambridge Philosophical Society* 75:405–435.
- Mulavwa M, Furuichi T, Yangozene K, et al. 2008. Seasonal changes in fruit production and party size of bonobos at Wamba. In: Furuichi T, Thompson J, editors. *The bonobos: behavior, ecology, and conservation*. New York: Springer. p 121–134.
- Myers Thompson JA. 2002. Bonobos of the lukuru wildlife research project. In: Boesch C, Hohmann G, Marchant LF, editors. *Behavioural diversity in chimpanzees and bonobos*. New York: Cambridge University Press. p 61–70.
- Newton-Fisher NE. 1999. Association by male chimpanzees: a social tactic? *Behaviour* 136:705–730.
- Newton-Fisher NE, Reynolds V, Plumtre AJ. 2000. Food supply and chimpanzee (*Pan troglodytes schweinfurthii*) party size in the Budongo Forest Reserve, Uganda. *International Journal of Primatology* 21:613–628.
- Nishida T. 1968. The social group of wild chimpanzees in the Mahali Mountains. *Primates* 9:167–224.

- Nishida T. 1979. The social structure of chimpanzees of the Mahale Mountains. In: Hamburg DA, McCown ER, editors. *The great apes*. Menlo Park: Benjamin/Cummings. p 73–121.
- Nishida T. 1990. *The chimpanzees of the Mahale mountains*. Tokyo: University of Tokyo Press.
- Nishida T, Hiraiwa-Hasegawa M. 1987. Chimpanzees and bonobos: cooperative relationships among males. In: Smuts BB, Cheney DL, Seyfarth RM, Wrangham RW, Struhsaker TT, editors. *Primate societies*. Chicago: University of Chicago Press. p 165–177.
- Nunes A. 1998. Diet and feeding ecology of *Ateles belzebuth belzebuth* at Maracá Ecological Station, Roraima, Brazil. *Folia Primatologica* 69:61–76.
- Nunn CL, Heymann EW. 2005. Malaria infection and host behavior: a comparative study of Neotropical primates. *Behavioral Ecology and Sociobiology* 59:30–37.
- Overdorff DJ. 1993. Similarities, differences, and seasonal patterns in the diets of *Eulemur rubriventer* and *Eulemur fulvus rufus* in the Ranomafana National Park, Madagascar. *International Journal of Primatology* 14:721–753.
- Overdorff D, Erhart E, Mutschler T. 2003. Fission–fusion in *Eulemur fulvus rufus* in southeastern Madagascar. *American Journal of Physical Anthropology* 38:92.
- Pereira ME, Seeligson ML, Macedonia JM. 1988. The behavioral repertoire of the black-and-white ruffed lemur, *Varecia variegata variegata* (Primates: Lemuridae). *Folia Primatologica* 51:1–32.
- Petter JJ, Albignac R, Rumpler Y. 1977. Mammifères lémurins (Primates, Prosimiens). *Faune de Madagascar*. Paris: ORSTROM/CNRS. 44. p 1–513.
- Pollock JI. 1979. Spatial distribution and ranging behavior in lemurs. In: Doyle GA, Martin RD, editors. *The study of prosimian behavior*. New York: Academic. p 359–409.
- Potts KB, Watts DP, Wrangham RW. 2011. Comparative feeding ecology of two communities of chimpanzees (*Pan troglodytes*) in Kibale National Park, Uganda. *International Journal of Primatology* 32:669–690.
- Pozo Rivera WE. 2001. Social behavior and diet of the spider monkey, *Ateles belzebuth*, in the Yasuni National Park, Ecuador. *Neotropical Primates* 9:74.
- Pozo Rivera WE. 2004. Agrupación y dieta de *Ateles belzebuth belzebuth* en el Parque Nacional Yasuni, Ecuador. *Anuar Investig Cientif* 2:77–102.
- Pride RE. 2005. Optimal group size and seasonal stress in ring-tailed lemurs (*Lemur catta*). *Behavioral Ecology* 16:550–560.
- Pruetz JD, Bertolani P. 2009. Chimpanzee (*Pan troglodytes verus*) Behavioral responses to stresses associated with living in a savanna-mosaic environment: implications for hominin adaptations to open habitats. *PaleoAnthropology* 252–262.
- Pusey A, Murray C, Wallauer W, et al. 2008. Severe aggression among female *Pan troglodytes schweinfurthii* at Gombe National Park, Tanzania. *International Journal of Primatology* 29:949–973.
- R Development Core Team. 2014. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing.
- Ramos-Fernandez G, Ayala-Orozco B. 2003. Population size and habitat use of spider monkeys at Punta Laguna, Mexico. In: Marsh LK, editor. *Primates in fragments: ecology and conservation*. New York, NY: Kluwer Academic: Plenum Publishers. p 191–209.
- Rasmussen DT. 1985. A comparative study of breeding seasonality and litter size in eleven taxa of captive lemurs (Lemur and Varecia). *International Journal of Primatology* 6:501–517.
- Ratsimbazafy J. 2002. On the brink of extinction and the process of recovery: responses of black-and-white ruffed lemurs (*Varecia variegata variegata*) to disturbance in Manombo [dissertation]. Stony Brook: Stony Brook University.
- Razakamaharavo VR, McGuire SM, Vasey N, Louis EE, Jr., Brenneman RA. 2010. Genetic architecture of two red ruffed lemur (*Varecia rubra*) populations of Masoala National Park. *Primates* 51:53–61.
- Ren B, Li D, Garber PA, Li M. 2012. Fission–fusion behavior in Yunnan snub-nosed monkeys (*Rhinopithecus bieti*) in Yunnan, China. *International Journal of Primatology* 33:1096–1109.
- Reynolds V. 2005. *The chimpanzees of the Budongo forest: ecology, behaviour, and conservation*. Oxford: Oxford University Press. p 297.
- Rifkin JL, Nunn CL, Garamszegi LZ. 2012. Do animals living in larger groups experience greater parasitism? A meta-analysis. *American Naturalist* 180:70–82.
- Rigamonti M. 1993. Home range and diet in red ruffed lemurs (*Varecia variegata rubra*) on the Masoala Peninsula, Madagascar. In: Ganzhorn J, Kappeler PM, editors. *Lemur social systems and their ecological basis*. New York: Plenum Press. p 25–39.
- Rodrigues de M, Carvalho O, Strier KB. 1998. Population variation in patch and party size in Muriquis (*Brachyteles arachnoides*). *International Journal of Primatology* 19:325–337.
- Sakura O. 1994. Factors affecting party size and composition of chimpanzees (*Pan troglodytes verus*) Bossou, Guinea. *International Journal of Primatology* 15:167–183.
- Schnell GD, Watt DJ, Douglas ME. 1985. Statistical comparison of proximity matrices: applications in animal behaviour. *Animal Behaviour* 33:239–253.
- Shimooka Y. 2003. Seasonal variation in association patterns of wild spider monkeys (*Ateles belzebuth belzebuth*) at La Macarena, Colombia. *Primates* 44:83–90.
- Slater KY, Schaffner CM, Aureli F. 2009. Sex differences in the social behavior of wild spider monkeys (*Ateles geoffroyi yucatanensis*). *American Journal of Primatology* 71:21–29.
- Snaith TV, Chapman CA, Rothman JM, Wasserman MD. 2008. Bigger groups have fewer parasites and similar cortisol levels: a multi-group analysis in red colobus monkeys. *American Journal of Primatology* 70:1–9.
- Snyder-Mackler N, Beehner JC, Bergman TJ. 2012. Defining higher levels in the multilevel societies of geladas (*Theropithecus gelada*). *International Journal of Primatology* 33:1054–1068.
- Stanford CB, Wallis J, Mpongo E, Goodall J. 1994. Hunting decisions in wild chimpanzees. *Behaviour* 131:1–18.
- Stanford CB. 1998. Predation and male bonds in primate societies. *Behaviour* 135:513–533.
- Sterck EH, Watts DP, van Schaik CP. 1997. The evolution of female social relationships in nonhuman primates. *Behavioral Ecology and Sociobiology* 41:291–309.
- Stevenson PR, Quinones MJ, Ahumada JA. 1998. Effects of fruit patch availability on feeding subgroup size and spacing patterns in four primate species at Tinigua National Park, Colombia. *International Journal of Primatology* 1:313–324.
- Strier KB. 1994. Myth of the typical primate. *Yearbook of Physical Anthropology* 37:233–271.
- Struhsaker TT, Leland L. 1979. Socioecology of five sympatric monkey species in the Kibale Forest, Uganda. *Advances in the Study of Behavior* 9:159–228.
- Stumpf R. 2011. Chimpanzees and bonobos: diversity within and between species. In: Campbell CJ, Fuentes A, MacKinnon KC, Bearder SK, Stumpf RM, editors. *Primates in perspective*. Oxford: Oxford University Press. p 321–344.
- Suarez S. 2003. Spatio-temporal foraging skills of white-bellied spider monkeys (*Ateles belzebuth belzebuth*) in the Yasuni National Park, Ecuador [dissertation]. Stony Brook: State University of New York.
- Sugiyama Y. 1968. Social organization of chimpanzees in the Budongo forest, Uganda. *Primates* 9:225–258.

- Sugiyama Y. 1981. Observations on the population dynamics and behavior of wild chimpanzees at Bossou, Guinea, 1979–1980. *Primates* 22:435–444.
- Sugiyama Y. 1988. Grooming interactions among adult chimpanzees at Bossou, Guinea, with special reference to social structure. *International Journal of Primatology* 9:393–407.
- Sugiyama Y, Koman J. 1979. Social structure and dynamics of wild chimpanzees at Bossou, Guinea. *Primates* 20:323–339.
- Sussman RW, Garber PA. 2011. Cooperation and competition in primate social interactions. In: Campbell CJ, Fuentes A, MacKinnon KC, Bearder SK, Stumpf RM, editors. *Primates in perspective*. Oxford: Oxford University Press. p 636–651.
- Swedell L, Plummer T. 2012. A papionin multilevel society as a model for hominin social evolution. *International Journal of Primatology* 33:1165–1193.
- Symington MM. 1987. Ecological and social correlates of party size in the black spider monkeys, *Ateles paniscus chamek* [dissertation]. Princeton, NJ: Princeton University.
- Symington MM. 1988a. Food competition and foraging party size in the black spider monkey (*Ateles paniscus chamek*). *Behaviour* 105:117–134.
- Symington MM. 1988b. Demography, ranging patterns, and activity budgets of black spider monkeys (*Ateles paniscus chamek*) in the Manu National Park, Peru. *American Journal of Primatology* 15:45–67.
- Symington MM. 1990. Fission–fusion social organization in *Ateles* and *Pan*. *International Journal of Primatology* 11:47–61.
- Tecot SR, Baden AL, Romine N, Kamilar J. 2013. Reproductive strategies and infant care in the Malagasy primates. In: Clancy K, Hinde K, Rutherford J, editors. *Primate developmental trajectories in proximate and ultimate perspectives*. Springer developments in primatology: progress and prospects book series. New York: Springer. p 321–359.
- Terborgh J, Janson CH. 1986. The socioecology of primate groups. *Annual Review of Ecological Systems* 17:1111–1135.
- Toborowsky C. 2008. Preliminary evidence of fission–fusion sub-grouping behavior in white-fronted brown lemurs (*Eulemur fulvus albifrons*) in Betampona Natural Reserve, Madagascar. *American Journal of Physical Anthropology* 46:208.
- Tutin CEG, McGrew WC, Baldwin PJ. 1983. Social organization of savanna-dwelling chimpanzees, *Pan troglodytes verus*, at Mt. Assirik, Senegal. *Primates* 24:154–173.
- Uehara S. 1981. The social unit of wild chimpanzees: a reconsideration based on the diachronic data accumulated at Kasoje in the Mahale Mountains, Tanzania. *Journal of African Studies* 20:15–32.
- van Hooft JARAM, van Schaik CP. 1994. Male bonds: affiliative relationships among nonhuman primate males. *Behaviour* 130:309–337.
- van Schaik CP. 1983. Why are diurnal primates living in groups? *Behaviour* 87:120–144.
- van Schaik CP. 1989. The ecology of social relationships amongst female primates. In: Standen V, Foley R, editors. *Comparative socioecology: the behavioural ecology of humans and other mammals*. Oxford: Blackwells Scientific Publications. p 195–218.
- van Schaik CP, van Hooft JARAM. 1983. On the ultimate causes of primate social systems. *Behaviour* 85:91–117.
- Vasey N. 2000. Niche separation in *Varecia variegata rubra* and *Eulemur fulvus albifrons*: I. Interspecific patterns. *American Journal of Physical Anthropology* 112:411–431.
- Vasey N. 2003. *Varecia*, ruffed lemurs. In: Goodman SM, Benstead J, editors. *The natural history of Madagascar*. Chicago: The University of Chicago Press. p 1332–1336.
- Vasey N. 2006. Impact of seasonality and reproduction on social structure, ranging patterns and fission–fusion social organization in red ruffed lemurs. In: Gould L, Sauther ML, editors. *Lemurs: ecology and adaptation*. New York: Springer/Kluwer. p 275–305.
- Vasey N. 2007. The breeding system of wild red ruffed lemurs (*Varecia rubra*): a preliminary report. *Primates* 48:41–54.
- Veiga LM, Pinto LP, Ferrari SF. 2006. Fission–fusion sociality in bearded sakis (*Chiropotes albinasus* and *Chiropotes satanas*) in Brazilian Amazonia. *International Journal of Primatology* 27:224.
- Vigilant L, Hofreiter M, Siedel H, Boesch C. 2001. Paternity and relatedness in wild chimpanzee communities. *Proceedings of the National Academy of Sciences* 98:12890–12895.
- Wakefield ML. 2008. Grouping patterns and competition among female *Pan troglodytes schweinfurthii* at Ngogo, Kibale National Park, Uganda. *International Journal of Primatology* 29:907–929.
- Wakefield ML. 2013. Social dynamics among females and their influence on social structure in an East African chimpanzee community. *Animal Behaviour* 85:1303–1313.
- Wallace RB. 1998. The behavioural ecology of black spider monkeys in north-eastern Bolivia [Ph.D. thesis]. Liverpool: University of Liverpool. p 282.
- Wallace RB. 2008. Towing the party line: territoriality, risky boundaries and male group size in spider monkey fission–fusion societies. *American Journal of Primatology* 70:271–281.
- White FJ. 1988. Party composition and dynamics in *Pan paniscus*. *International Journal of Primatology* 9:179–193.
- White FJ. 1989. The social organization of pygmy chimpanzees. In: Heltne PG, Marquardt LA, editors. *Understanding chimpanzees*. Massachusetts: Harvard University Press. p 194–207.
- White FJ. 1991. Social organization, feeding ecology, and reproductive strategy of ruffed lemurs, *Varecia variegata*. In: Ehara A, Kimura T, Takenaka O, Iwamoto M, editors. *Primateology today*. New York: Elsevier Science Publishers. p 278.
- White FJ. 1992. Pygmy chimpanzee social organization: variation with party size and between study sites. *American Journal of Primatology* 26:203–214.
- White FJ, Wrangham RW. 1988. Feeding competition and patch size in the chimpanzee species *Pan paniscus* and *Pan troglodytes*. *Behaviour* 105:148–164.
- Whitehead H. 2008. *Analyzing animal societies: quantitative methods for vertebrate social analysis*. Chicago: University of Chicago Press.
- Whitehead H. 2009. SOCPROG programs: analyzing animal social structures. *Behavioral Ecology and Sociobiology* 63:765–778.
- Williams JM, Pusey AE, Carlis JV, Farm BP, Goodall J. 2002. Female competition and male territorial behaviour influence female chimpanzees' ranging patterns. *Animal Behaviour* 63:347–360.
- Wittiger L, Boesch C. 2013. Female gregariousness in Western Chimpanzees (*Pan troglodytes verus*) is influenced by resource aggregation and the number of females in estrus. *Behavioral Ecology and Sociobiology* 67:1097–1111.
- Wrangham RW. 1977. Feeding behaviour of chimpanzees in Gombe National Park, Tanzania. In: Clutton-Brock TH, editor. *Primate ecology*. London: Academic Press. p 504–538.
- Wrangham RW. 1980. An ecological model of female-bonded primate groups. *Behaviour* 75:262–300.
- Wrangham RW. 1986. Ecology and social relationships in two species of chimpanzee. In: Rubenstein DI, Wrangham RW, editors. *Ecology and social evolution: birds and mammals*. Princeton, USA: Princeton University Press. p 352–378.
- Wrangham RW. 2000. Why are male chimpanzees more gregarious than mothers? A scramble competition hypothesis. In: Kappeler PM, editor. *Primate ecology: studies of feeding and ranging behavior in lemurs, monkeys and apes*. Cambridge: Cambridge University Press. p 248–258.

- Wrangham R, Smuts B. 1980. Sex differences in the behavioural ecology of chimpanzees in the Gombe National Park, Tanzania. *Journal of Reproduction and Fertility Supplement* 28:13–31.
- Wrangham RW, Clark AP, Isabirye-Basuta G. 1992. Female social relationships and social organization of Kibale forest chimpanzees. In: Nishida T, McGrew WC, Marler P, Pickford M, de Waal FB, editors. *Topics in primatology*. Tokyo: University of Tokyo Press. p 81–98.
- Wrangham RW, Chapman CA, Clark-Arcadi AP, Isabirye-Basuta G. 1996. Social ecology of Kanyawara chimpanzees: implications for understanding the costs of great apes groups. In: McGrew WC, Marchant LF, Nishida T, editors. *Great ape societies*. Cambridge: Cambridge University Press. p 45–57.
- Wright PC. 1999. Lemur traits and Madagascar ecology: coping with an island environment. *Yearbook of Physical Anthropology* 42:31–72.
- Wright PC, Erhart EM, Tecot S, et al. 2012. Long-term lemur research at centre Valbio, Ranomafana National Park, Madagascar. In: Kappeler PM, Watts DP, editors. *Long-term field studies of primates*. New York, NY: Springer. p 67–100.